

PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁷ : C12Q		A2	(11) International Publication Number: WO 00/46391
			(43) International Publication Date: 10 August 2000 (10.08.00)
(21) International Application Number: PCT/US00/02920 (22) International Filing Date: 2 February 2000 (02.02.00) (30) Priority Data: 09/243,277 2 February 1999 (02.02.99) US (63) Related by Continuation (CON) or Continuation-in-Part (CIP) to Earlier Application US 09/243,277 (CIP) Filed on 2 February 1999 (02.02.99)		(81) Designated States: AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CR, CU, CZ, DE, DK, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG). Published <i>Without international search report and to be republished upon receipt of that report.</i>	
(71) Applicant (for all designated States except US): THE TRUSTEES OF COLUMBIA UNIVERSITY IN THE CITY OF NEW YORK [US/US]; West 116th Street and Broadway, New York, NY 10027 (US). (72) Inventors; and (75) Inventors/Applicants (for US only): FISHER, Paul, B. [US/US]; 15 Gordon Place, Scarsdale, NY 10583 (US). LESZCYNIECKA, Magdalena [US/US]; 453 Westend Avenue, Clifton, NJ 07011 (US). (74) Agent: WHITE, John, P.; Cooper & Dunham LLP, 1185 Avenue of the Americas, New York, NY 10036 (US).			
(54) Title: GENES DISPLAYING ENHANCED EXPRESSION DURING CELLULAR SENESCENCE AND TERMINAL CELL DIFFERENTIATION AND USES THEREOF			
(57) Abstract This invention provides isolated nucleic acid molecules encoding an OLD-35 protein, OLD-64 protein, OLD-137 protein, OLD-139 protein, OLD-142 protein or OLD-175 protein. This invention further provides a purified OLD-35 protein, OLD-64, OLD-137, OLD-139, OLD-142 and OLD-175. Finally, this invention provides different uses of the nucleic acids and proteins.			

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece			TR	Turkey
BG	Bulgaria	HU	Hungary	ML	Mali	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MN	Mongolia	UA	Ukraine
BR	Brazil	IL	Israel	MR	Mauritania	UG	Uganda
BY	Belarus	IS	Iceland	MW	Malawi	US	United States of America
CA	Canada	IT	Italy	MX	Mexico	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NE	Niger	VN	Viet Nam
CG	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NO	Norway	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	NZ	New Zealand		
CM	Cameroon			PL	Poland		
CN	China	KR	Republic of Korea	PT	Portugal		
CU	Cuba	KZ	Kazakhstan	RO	Romania		
CZ	Czech Republic	LC	Saint Lucia	RU	Russian Federation		
DE	Germany	LI	Liechtenstein	SD	Sudan		
DK	Denmark	LK	Sri Lanka	SE	Sweden		
EE	Estonia	LR	Liberia	SG	Singapore		

GENES DISPLAYING ENHANCED EXPRESSION DURING
CELLULAR SENESENCE AND TERMINAL CELL

5 DIFFERENTIATION AND USES THEREOF

This application claims priority and is a continuation-in-part application of U.S. Serial No. 09/243,277, filed February 2, 1999, the contents of which is hereby
10 incorporated by reference.

Throughout this application, various publications are referred to by arabic numeral within parentheses. Full citations for these publications are presented immediately
15 before the claims. Disclosures of these publications in their entireties are hereby incorporated by reference into this application in order to more fully describe the state of the art to which this invention pertains.

20 BACKGROUND OF THE INVENTION

Normal cells cultured *in vitro* lose their proliferative potential after a finite number of doublings in a process described as cellular senescence (Hayflick and Moorehead, 1976). This phenomenon is not only well-established in human
25 diploid fibroblasts based on the studies of Hayflick and Moorehead (1976), but it has also been validated by investigations using many additional cell types (Goldstein et al., 1990; Murano et al., 1991). These investigations document an inverse correlation between replicative
30 senescence and donor age and a direct relationship between replicative senescence and donor species lifespan (Hayflick and Moorehead, 1976; Goldstein et al., 1990; Murano et al., 1991). In agreement with this association, cells from patients with premature aging syndromes, such as Werner's
35 syndrome and Progeria, achieve a quiescent state much more rapidly than normal human fibroblasts. In this context, if similar senescence related changes occur in normal fibroblasts, albeit with delayed kinetics, these cell systems represent excellent models for studying senescence *in vitro*
40 and identifying genes relevant to the aging process.

-2-

Senescence is characterized by changes in cell morphology, lack of responsiveness to mitogenic stimulation and irreversible growth arrest. However, cells can withdraw from the cell cycle and become non-dividing not only during
5 senescence but also during the processes of DNA damage, apoptosis or terminal differentiation. While senescence is a time-dependent process (Campisi et al., 1995), terminal differentiation can be induced in a variety of cell types by appropriate treatment (Roberts et al., 1999). For example,
10 terminal differentiation can be induced by cAMP treatment in melanocytes (Medrano et al., 1994). Gene expression analysis in terminally differentiated versus senescent melanocytes indicates both similarities and differences (Medrano et al., 1994). Although both pathways result in an elevation in p21
15 and an inability to phosphorylate ERK2, only the differentiated cells display elevated levels of p27 and the melanocyte-specific transcription factor (MITF) (Medrano et al., 1994; Smith and Pereira-Smith, 1996).

20 Human melanoma represents an excellent model for studying irreversible growth arrest and terminal differentiation, since these physiological changes can be chemically induced by IFN- β plus mezerein (MEZ) (Fisher et al., 1985; Jiang et al., 1994a). The induction of terminal differentiation in
25 HO-1 human melanoma cells correlates with up-regulation of c-jun, jun-B, α_5 Integrin, β_1 Integrin, fibronectin, HLA Class I, ISG-54, ISG-15 and gro/MSGA as well as down-regulation of c-myc (Jiang et al., 1993a). To define the repertoire of genes differentially expressed during induction of
30 irreversible growth arrest and terminal differentiation in human melanoma cells we have used a rapid and efficient differentiation induction subtraction hybridization (DISH) approach (Jiang and Fisher, 1993). Using this approach alone and in combination with high throughput screening strategies,
35 microchip DNA arrays, a large number of novel genes of potential relevance to growth control and terminal differentiation have been identified and cloned (Jiang et

-3-

al., 1995a, 1995b; Lin et al., 1996, 1998; Huang et al., 1999).

On the basis of the considerations described above, it is probable that specific differentially expressed genes may be present within a terminally differentiated cDNA library that also display modified expression during cellular senescence. To begin to identify these overlapping genes, a temporally spaced subtracted differentiation inducer treated H0-1 human melanoma library was screened with RNA isolated from senescent human fibroblasts. Such a screening protocol yielded twenty-eight known and ten novel cDNAs. Subsequent Northern and reverse Northern blotting analyses revealed differential expression of specific cDNAs. Expression of one of these cDNAs, Old-35 was restricted to terminal differentiation and senescence. In this context, this gene may contribute to pathways leading to growth arrest, a defining component of senescence and terminal differentiation.

Interferons (IFNs) comprise a family of related cytokines with diverse including antiviral, antiproliferative, antitumor and immunomodulatory activities (Stark et al., 1998; *Roberts et al., 1999). IFN studies have focused on two main areas; one involving the clinical use of IFN for therapeutic purposes (Guttermann, 1994), the other employing the IFN system as a paradigm to study the mammalian JAK/STAT signaling cascade (Darnell et al., 1994) that leads to IFN-stimulated gene (ISG) activation. To date, the most extensively studied ISGs include RNA-activated protein kinase (PKR), the 2'-5' oligoadenylate synthetase and the MX proteins (Stark et al., 1998, *Der et al., 1998).

Post-transcriptional regulation of mRNA levels is a pivotal control point in gene expression. Early response genes, such as cytokines, lymphokines and proto-oncogenes are regulated by a cis-acting adenylate-uridylate-rich element (ARE) found in the 3' untranslated region (UTR) of the mRNA (Caput et

-4-

al., 1986; Shaw and Kamen, 1988; Chen and Shyu, 1995; Myer et al., 1997). Currently, three classes of destabilizing elements have been identified: AUUUA-lacking elements and AUUUA-containing elements grouped into those with scattered 5 AUUUA motifs (such as proto-oncogenes) and those with overlapping AUUUA motifs (such as growth factors) (Chen et al., 1995; Myer et al., 1997). A transfer of 3'UTR containing ARE to 3'UTR of a stable message, such as β -globin, targets this very stable mRNA for rapid degradation (Shaw and Kamen, 10 1988). In contrast, the removal of an ARE stabilizes an otherwise labile message (*Miller et al., 1984; *Lee et al., 1988).

The present studies describe the cloning and initial 15 characterization of a novel gene, Old-35, from a terminally differentiated human melanoma cDNA library. mRNA stability studies document that Old-35 mRNA, which contains ARE elements, may be stabilized in HO-1 cells by treatment with IFN- β and IFN- β + MEZ. Based on the growth suppressive effect 20 of IFN- β on HO-1 cells, as well as the increased stability of Old-35 during confluence and senescence, it is possible that this gene plays a prominent role in growth suppression induced by this cytokine. Further experimentation is required to define the precise role of Old-35 in IFN signaling, 25 terminal differentiation and cellular senescence. Full-length cloning and subsequent protein analyses should provide insights into the function of this potentially important gene in the processes of aging and differentiation.

30 Since the processes of terminal differentiation and senescence exhibit strikingly similar characteristics, it is possible that related and overlapping genes and gene expression changes associate with and mediate both of these phenomena. Old-35 was isolated by screening a subtracted 35 human melanoma cDNA library enriched for genes related to growth arrest and terminal differentiation with RNA from senescent human fibroblasts. This cDNA encodes an IFN- β inducible gene expressed during different types of growth

-5-

arrest including confluence, senescence and terminal differentiation. Old-35 RNA exhibits increased stability in IFN- β and INF- β + MEZ treated H0-1 human melanoma cells. Steady-state mRNA for Old-35 is also highly expressed in 5 heart and brain, human tissues without active regenerative properties. Judging from the pattern of Old-35 expression, it is possible that this gene may play a prominent role during growth arrest and in this context contributes to the important processes of senescence and terminal 10 differentiation.

-6-

SUMMARY OF THE INVENTION

This invention provides isolated nucleic acid molecule encoding an old 35 protein, 64 protein, 137 protein, 139 protein, 142 protein and a 175 protein. The isolated nucleic acid may be a DNA, genomic DNA, cDNA, synthetic DNA or RNA. The isolated nucleic acid has a sequence substantially the same as SEQ ID. Nos. 39, 19, 31, 32, 34 and 38 which are respectively Old 35, old 64, old 137, old 139, old 142 and old 175.

10

This invention also provides a nucleic acid molecule of at least 15 nucleotides capable of specifically hybridizing with a sequence included within the sequence of a nucleic acid molecule encoding an old 35 protein, 64 protein, 137 protein, 139 protein, 142 protein and a 175 protein. The nucleic acid probe may be DNA, genomic DNA, cDNA, synthetic DNA or RNA.

This invention further provides a host vector system for the production of a protein having the biological activity of old 35, 64, 137, 139, 142 and 175. The isolated old 35, 64, 137, 139, 142 and 175 nucleic acid molecule is linked to a promoter of RNA transcription and then to a plasmid. The suitable host is a bacterial cell, insect cell, or animal cell, depending on the type of promoter and plasmid used. This invention also provides a method of producing a protein having the biological activity of old 35, 64, 137, 139, 142 and 175, which comprises growing the selected host vector system under suitable conditions permitting production of the protein and recovering the protein so produced.

30

This invention further provides purified protein of old 35, 64, 137, 139, 142 and 175. Such purified old 35, 64, 137, 139, 142 and 175 will be useful for inhibiting growth of cancer cells. This invention provides a method of contacting the cancer cells with an amount of old 35, 64, 137, 139, 142 and 175 at a concentration effective to inhibit growth of cancer cells. This invention further provides a method of determining whether a cell is senescent by (a)isolating the

-7-

nucleic acids in the cell (b) hybridizing the isolated nucleic acids with the nucleic acid of old 35 or 64 under conditions permitting hybrids formation and (c) detecting the expression of old 35 or old 64 in the cell. This invention
5 further provides a method of determining whether a cell has growth arrest by (a) isolating the nucleic acids in the cell; (b) hybridizing the isolated nucleic acids with the nucleic acid of old 35 or 64 under conditions permitting hybrids formation; and (c) detecting the expression of old 35 or old
10 64 in the cell. This invention further provides a method of determining whether a cell has terminal differentiation by (a) isolating the nucleic acids in the cell; (b) hybridizing the isolated nucleic acids with the nucleic acid of old 35 or 64 under conditions permitting hybrids formation; and (c)
15 detecting the expression of old 35 or old 64 in the cell. Further, this invention provides that the detector used is a DNA, RNA or protein. This invention also provides a method of regenerating tissue with an inhibitor of old 35 protein at a concentration effective to regenerate said tissues.
20 This invention provides a method of anti-aging in a cell comprising contacting the cell with an agent for inhibiting expression of old 35 at a concentration effective to reverse growth arrest in the cell. Finally, this invention provides a pharmaceutical composition for stimulating cell growth
25 comprising a pharmaceutically acceptable carrier and purified old 35 or old 64 at a concentration effective to stimulate cell growth.

BRIEF DESCRIPTION OF FIGURES

- Figure 1** Expression of Old-35 in H0-1 human melanoma cells treated with IFN- β or FN- β + MEZ, young human fibroblasts and two different types of senescent Progeria human fibroblasts. Northern blot contains 10 μ g of total RNA from control untreated H0-1 (lane 1), IFN- β treated (2,000 U/ml) H0-1 (lane 2), IFN- β + MEZ treated (2,00U/ml + 10ng/ml) H0-1 (lane 3), young fibroblasts (GM01379) (lane 4), and two senescent Progeria cell lines (AG01976) (lane 5) (AG0989B) (lane 6). Blots were exposed for autoradiography for 1, 4 or 24 hr. EtBr staining for quantification of gel loading and determining RNA quality.
- Figure 2** Effect of IFN- α , IFN- β , IFN- γ , TNF- α and IFN- β + MEZ on Old-35 expression in H0-1 cells. All Northern blots contain 10 mg of total RNA. (A) Time course induction of Old-35 by IFN- β in H0-1 cells. Cells were seeded at ~60% confluence and treated with IFN- β (2,000 units/ml) and RNA was isolated at the indicated time. U = RNA from control, untreated cells. (B) Dose response expression of Old-35 in H0-1 cells treated with IFN- β (2,000 units/ml). RNAs were isolated after 24 hr treatment. (C) Effect of IFN- α (I α), IFN- β (I β), IFN- γ (I γ) and TNF- α (T α) on Old-35 expression in H0-1 cells. RNAs were isolated after 15 hr treatment with 1,000 units/ml of the different agents. U = RNA from control, untreated cells. (D) Time course induction of Old-35 by IFN- β + MEZ in H0-1 cells. RNAs were isolated from cells treated with 2,000 units/ml of IFN- β + 10 ng/ml of MEZ.
- Figure 3** Expression of Old-35 in various human tissues and during mouse development. (A) Northern blot

-9-

contains 2 μ g of poly A⁺ RNA per lane from eight different human tissues. Lanes 1-8 contain, in order, RNA from human heart, brain, placenta, lung, liver, skeletal muscle, kidney, and pancreas (Clontech). (B) Northern blot contains 10 μ g of total RNA from mouse embryos. The number of days signifies days post-gestation.

Figure 4 Sequence comparison between human and the mouse homologue of Old-35. Upper panel sequence of human Old-35 (h-Old-35); Middle panel: sequence of mouse Old-35 (m-Old-35); and Lower panel: shared consensus sequences between human and mouse Old-35.

Figure 5 Expression of Old-35 in IDH4 cells grown in the presence or absence of Dex. Northern blot contains 10 μ g of total RNA per lane from IDH-4 cells. + Dex = cells grown continuously in the presence of 10⁻⁶ M Dex; - Dex = cells grown for the indicated days in the absence of Dex. For the latter experiment, cells were grown in the presence of Dex and then shifted to charcoal stripped media and grown for 3, 5, 7 and 14 days without Dex.

Figure 6 Expression of Old-35 and p21 during cell cycle progression in human skin fibroblasts. Northern blot contains 10 μ g of total RNA per lane from normal human fibroblasts. Confluent normal fibroblasts (C) were trypsinized and reseeded (1:2). Total RNA was collected at 5, 15 and 20 hr after reseeding. At 20hr following subculture, the cells were 90% confluent.

Figure 7 AU rich sequences found in the 3' untranslated region (UTR) of several lymphokine and

-10-

protooncogene mRNAs. Abbreviations:
Abbreviations: Hu-human, GM-CSF =
granulocyte-monocyte colony stimulating factor;
IFN- α = interferon α ; IL 2 = Interleukin 2; TNF
5 = tumor necrosis factor; c-fos = fos
proto-oncogene. The underlined/overlined AUUUA
motif is the largest sequence common to all mRNAs
shown. References: HuGM-CSF (Wong et al., 1985),
HuIFN- α (Goeddel et al., 1983), Hu IL 2 (Kashima
10 et al., 1985), HuTNF (Nedwin et al., 1985), Hu
c-fos (van Straaten et al., 1983).

Figure 8 Effect of cycloheximide treatment on Old-35
expression in H0-1 cells and the half-life of
15 Old-35 mRNA in IFN- β + MEZ treated H0-1 cells.
Each lane in the Northern blots contains 10 μ g of
total RNA. (A) H0-1 cells were pre-treated with
cyclohexamide 50mg/ml for 30 min and then treated
with IFN- β for 2, 3 or 4 hr (lanes 2, 3, and 4,
20 respectively). H0-1 cells were pre-treated with
IFN- β for 5 hr (lane 5) and then treated with
cycloheximide for 15 hr (lane 6). U = RNA from
control untreated H0-1 cells. (B) Half-life of
Old-35 mRNA in IFN- β + MEZ (IM) (2,000 units/ml
25 + 10 ng/ml) treated H0-1 cells. Cells were
incubated with IM for 15 hr and then exposed to
ActD (50 mg/ml) for 2, 6, 8, 10 and 12 hr. U =
RNA from control untreated H0-1 cells. AD = RNA
from control H0-1 cells treated with ActD (5
30 μ g/ml).

Figure 9 DNA sequence and predicted encoded protein of
Old-35. (A) cDNA sequence of Old-35. Alternate
polyadenylation site is underlined. This site is
35 present in 10% of all cDNAs (*Manley et al.,
1988). (B) Predicted protein encoded by the
Old-35 cDNA.

-11-

Figure 10 Sequence similarity between the bacterial protein PNPase and the predicted protein sequence of Old-35. Upper Panel: *Bacillus subtilis* PNPase sequence. Middle Panel: predicted human Old-35 protein sequence. Lower Panel: regions of consensus amino acids between the bacterial PNPase protein sequence and the predicted Old-35 protein sequence. Black boxed areas indicates amino acid identity and gray boxed areas indicate amino acid similarities between the bacterial PNPase and the predicted Old-35 encoded protein.

Figure 11 Northern Blot of HO-1, confluent HO-1, IFN- β treated, IFN- β +MEZ treated HO-1 treated with Actinomycin D (50mg/ml). Total RNAs were collected 2,4,6,8,10,12 after the AD treatment. Old-35 cDNA was used as a probe. Ethyidium Bromide was shown for loading control

Figure 12 Northern Blot of IDH4 and AR5 cells. IDH4 cells contain dexamethasone (DEX) inducible mouse mammary tumor virus-driven simian virus 40 T-antigen. Total RNA was extracted from cells treated with DEX (indicated as +), and from cells growing without DEX for 3,5,7, and 14 days). AR5 cells contain temperature sensitive simian virus 40 T-antigen. Total RNA was collected from cells at 35C and 1,3,7,14 days after shift to 39C. Old-35 and p21 were used as a probe.

Figure 13 Structure of Old-35 gene. RnasePH, KH, S1 signify domains found in Old-35 cDNA. Top picture shows two different versions of Old-35 which vary in the 3'UTR length
The bottom picture shows cloning of the Old-35 cDNA using C-ORF technique.

-12-

Figure 14

Localization of GFP-Old-35, and GFP alone in HeLa cells.

Figure 15

5 *In situ* hybridization to mouse embryo (9.5 days) using murine Old-35. The arrows indicate the expression in the spinal column.

Figure 16

10 Northern blot of HO-1 cells treated with different subtypes of IFN- α using Old-35 as a probe. IFN- β was used as a control.

-13-

DETAILED DESCRIPTION OF THE INVENTION

In order to facilitate an understanding of the Experimental Details section which follows, certain frequently occurring methods and/or terms are described in Sambrook, et al. 5 (1989).

Throughout this application, the following standard abbreviations are used throughout the specification to indicate specific nucleotides:

10	C=cytosine	A=adenosine
	T=thymidine	G=guanosine

This invention provides an isolated nucleic acid molecule encoding an OLD-35 or OLD-64 protein. In an embodiment, the 15 above nucleic acid molecule comprises a nucleic acid having a sequence substantially the same as set forth in SEQ. ID. No.39 or 19.

This invention also provides isolated nucleic acid molecules 20 encoding an OLD-137, OLD-139, OLD-142, or OLD-175 protein. In an embodiment, the nucleic acid comprises a nucleic acid having a sequence substantially the same as set forth in SEQ. ID. Nos.31, 32, 34 or 38. The above-described nucleic acid may be DNA, genomic DNA, cDNA, synthetic DNA, or RNA.

25 This invention also encompasses nucleic acid which encode amino acid sequences which differ from those of OLD-35, OLD-64, OLD-137, OLD-139, OLD-142 or OLD-175, but which should not produce phenotypic changes. Alternatively, this 30 invention also encompasses DNAs and cDNAs which hybridize to the DNA and cDNA of the subject invention. Hybridization methods are well known to those of skill in the art.

The DNA molecules of the subject invention also include DNA 35 molecules coding for protein analogs, fragments or derivatives of antigenic proteins which differ from naturally-occurring forms in terms of the identity or location of one or more amino acid residues (deletion analogs

-14-

containing less than all of the residues specified for the protein, substitution analogs wherein one or more residues specified are replaced by other residues and addition analogs wherein one or more amino acid residues is added to a 5 terminal or medial portion of the proteins) and which share some or all properties of naturally-occurring forms. These sequences include: the incorporation of codons "preferred" for expression by selected non-mammalian host; the provision of sites for cleavage by restriction endonuclease enzymes; 10 and the provision of additional initial, terminal or intermediate DNA sequences that facilitate construction of readily expressed vectors.

The nucleic acid molecule described and claimed herein are 15 useful for the information which they provide concerning the amino acid sequence of the protein and as products for the large scale synthesis of the protein by a variety of recombinant techniques. The molecule is useful for generating new cloning and expression vectors, transformed 20 and transfected procaryotic and eukaryotic host cells, and new and useful methods for cultured growth of such host cells capable of expression of the protein and related products.

The invention also provides fragments or portion of the Old 25 gene or protein wherein the biological activity of said gene product is maintained. Such fragment or portion may join to other amino acid sequence to create a multi-functional molecule. It is within the ordinary skill to determine such biologically active fragment or portion. A trimming 30 experiment may be performed to define said fragment of portion.

Old-35, Old-64, Old-137, Old-139, Old-142 or Old-175 may be isolated in a variety of vertebrates. In an embodiment, a 35 human Old-35, Old-64, Old-137, Old-139, Old-142 and Old-175 are isolated.

The isolated nucleic molecule of Old-35, Old-64, Old-137,

-15-

Old-139, Old-142 and Old-175 are represented respectively by SEQ. ID. Nos. 39, 19, 31, 32, 34 and 38.

This invention provides a nucleic acid molecule of at least 5 15 nucleotides capable of specifically hybridizing with a sequence included within the sequence of a nucleic acid molecule encoding a Old-35, Old-64, Old-137, Old-139, Old-142 or Old-175. In an embodiment, the nucleic acid is DNA, genomic DNA, cDNA, synthetic DNA or RNA.

10

As used herein, the phrase "specifically hybridizing" means the ability of a nucleic acid molecule to recognize a nucleic acid sequence complementary to its own and to form double-helical segments through hydrogen bonding between 15 complementary base pairs. The nucleic acid molecule will be specific to said Old genes i.e. under appropriate conditions, the molecule will only hybridize with said old gene and no other genes. Said molecule may contain an unique sequence of said Old gene.

20

Nucleic acid probe technology is well known to those skilled in the art who will readily appreciate that such probes may vary greatly in length and may be labeled with a detectable label, such as a radioisotope or fluorescent dye, to 25 facilitate detection of the probe.

Probe molecules may be produced by insertion of a nucleic acid molecule which encodes OLD-35, OLD-64, OLD-137, OLD-139, OLD-142 or OLD-175 protein or a fragment thereof into 30 suitable vectors, such as plasmids or bacteriophages, followed by transforming into suitable bacterial host cells, replication in the transformed bacterial host cells and harvesting of the DNA probes, using methods well known in the art. Alternatively, probes may be generated chemically from 35 DNA synthesizers.

The probes are useful for 'in situ' hybridization to locate tissues which express this gene, or for other hybridization

-16-

assays for the presence of this gene or its mRNA in various biological tissues.

The invention also provides an antisense nucleic acid molecule comprising a sequence complementary to the nucleic acid which encodes OLD-35, OLD-64, OLD-137, OLD-139, OLD-142 or OLD-175 protein or a fragment thereof. In an embodiment, the antisense nucleic acid molecule is capable of inhibiting the expression of the hybridized gene.

10

This invention also provides the above-described isolated nucleic acid molecule operatively linked to a promoter of RNA transcription. This invention further provides a vector which comprises the above-described isolated nucleic acid molecule.

Vectors which comprise the isolated nucleic acid molecule described hereinabove also are provided. Suitable vectors comprise, but are not limited to, a plasmid or a virus. These vectors may be transformed into a suitable host cell to form a host cell vector system for the production of a protein having the biological activity of OLD-35, OLD-64, OLD-137, OLD-139, OLD-142 or OLD-175 protein or a fragment thereof.

25

This invention further provides an isolated DNA, genomic DNA, cDNA, synthetic DNA or RNA molecule described hereinabove wherein the host cell is selected from the group consisting of bacterial cells (such as E.coli), yeast cells, fungal cells, insect cells and animal cells. Suitable animal cells include, but are not limited to Vero cells, HeLa cells, Cos cells, CV1 cells and various primary mammalian cells.

This invention provides a purified, OLD-35 protein, a purified, OLD-64 protein, a purified, OLD-137 protein, a purified, OLD-139 protein, a purified, OLD-142 protein, and a purified, OLD-175 protein.

-17-

This invention also provides a protein encoded by the above-described isolated nucleic acid molecule.

This invention also provides an antibody or antigen-binding
5 fragment thereof that specifically binds to OLD-35, OLD-64, OLD-137, OLD-139, OLD-142 or OLD-175 protein. In an embodiment, the antibody is a monoclonal antibody.

Polyclonal antibodies against these proteins may be produced
10 by immunizing animals using selected peptides determined from the decoded amino acid sequences. Monoclonal antibodies are prepared using hybridoma technology by fusing antibody producing B cells from immunized animals with myeloma cells and selecting the resulting hybridoma cell line producing the
15 desired antibody. Alternatively, monoclonal antibodies may be produced by in vitro techniques known to a person of ordinary skill in the art. These antibodies are useful to detect the expression of the OLD proteins in living animals, in humans, or in biological tissues or fluids isolated from
20 animals or humans.

This invention provides a method of inhibiting growth of cancer cells comprising contacting the cancer cells with an amount of purified OLD-35, OLD-64 protein or a portion
25 thereof effective to inhibit growth of cancer cells.

This invention also provides a method for reversing the cancerous phenotype of a cancer cell which comprises introducing a nucleic acid comprising an Old-35 or Old-64
30 gene or a portion thereof into the cell under conditions permitting the expression of the gene so as to thereby reverse the cancerous phenotype of the cell.

This invention provides a method for reversing the cancerous
35 phenotype of a cancer cell in a subject which comprises introducing a nucleic acid molecule comprising an Old-35 or Old-64 gene or a portion thereof into the subject's cancerous cell under conditions permitting expression of the gene in

-18-

the subject's cell so as to thereby reverse the cancerous phenotype of the cell.

In an embodiment of the method, the nucleic acid molecule
5 comprises a vector. In a further embodiment, the Old-35 or
Old-64 gene is linked to a regulatory element such that its
expression is under the control of the regulatory element.
In a still further embodiment, the regulatory element is a
tissue specific regulatory element. In a still further
10 embodiment, the regulatory element is inducible or
constitutive. Inducible regulatory element like an inducible
promoter is known in the art. Regulatory element such as
promoter which can direct constitutive expression is also
known in the art.

15

In a separate embodiment, the regulatory element is a tissue
specific regulatory element. The expression of the inserted
gene will then be tissue-specific.

20 Methods to introduce a nucleic acid molecule into cells have
been well known in the art. Naked nucleic acid molecule may
be introduced into the cell by direct transformation.
Alternatively, the nucleic acid molecule may be embedded in
liposomes. Accordingly, this invention provides the above
25 methods wherein the nucleic acid is introduced into the cells
by naked DNA technology, adenovirus vector, adeno-associated
virus vector, Epstein-Barr virus vector, Herpes virus vector,
attenuated HIV vector, retroviral vectors, vaccinia virus
vector, liposomes, antibody-coated liposomes, mechanical or
30 electrical means. The above recited methods are merely
served as examples for feasible means of introduction of the
nucleic acid into cells. Other methods known may be also be
used in this invention.

35 This invention provides a method for reversing the cancerous
phenotype of a cancer cell which comprises introducing OLD-
35 or OLD-64 protein or a portion thereof into the cancerous
cell so as to thereby reverse the cancerous phenotype of the

-19-

cell.

This invention provides a method for reversing the cancerous phenotype of a cancer cell in a subject which comprises
5 introducing OLD-35 or OLD-64 protein into the subject's cancerous cell so as to thereby reverse the cancerous phenotype of the cell. In an embodiment, the cancer cell is a breast, cervical, colon, pancreatic, thyroid, skin, brain, prostate, nasopharyngeal, lung, glioblastoma multiforme,
10 lymphoma, leukemia, connective tissue, nervous system cell or basal cell.

This invention further provides a pharmaceutical composition which comprises an amount of a nucleic acid molecule
15 comprising Old-35, Old-64 gene or portion thereof effective to reverse the cancerous phenotype of a cancer cell and a pharmaceutically acceptable carrier. In an embodiment, the nucleic acid molecule comprises a vector. In a further embodiment, the vector is an adenovirus vector, adeno-
20 associated virus vector, Epstein-Barr virus vector, Herpes virus vector, attenuated HIV vector, retrovirus vector or vaccinia virus vector.

This invention also provides a pharmaceutical composition
25 comprising an amount of OLD-35 or OLD-64 protein effective to reverse the cancerous phenotype of a cancer cell and a pharmaceutically acceptable carrier. In an embodiment, the cancer cell is a breast, cervical, colon, pancreatic, thyroid, skin, brain, prostate, nasopharyngeal, lung,
30 glioblastoma multiforme, lymphoma, leukemia, connective tissue, nervous system or basal cell.

In an embodiment of the above methods, the nucleic acid comprises a vector. The vector includes, but is not limited
35 to, an adenovirus vector, adeno-associated virus vector, Epstein-Barr virus vector, Herpes virus vector, attenuated HIV vector, retrovirus vector and vaccinia virus vector.

-20-

As used herein, the term "pharmaceutically acceptable carrier" encompasses any of the standard pharmaceutical carriers. The pharmaceutical composition may be constituted into any form suitable for the mode of administration
5 selected. Compositions suitable for oral administration include solid forms, such as pills, capsules, granules, tablets, and powders, and liquid forms, such as solutions, syrups, elixirs, and suspensions. Forms useful for parenteral administration include sterile solutions,
10 emulsions, and suspensions.

In the practice of the method administration may comprise daily, weekly, monthly, hourly or by peak and trough, the precise frequency being subject to various variables such
15 as age and condition of the subject, amount to be administered, half-life of the agent in the subject, area of the subject to which administration is desired and the like.

In connection with the method of this invention, a
20 therapeutically effective amount may include dosages which take into account the size and weight of the subject, the age of the subject, the severity of the symptom, the efficacy of the agent and the method of delivery of the agent. One of ordinary skill in the art would be readily able to determine
25 the exact dosages and exact times of administration based upon such factors.

This invention provides a method of determining whether a cell is senescent comprising measurement of the expression
30 of Old-35 gene, wherein the expression of the Old-35 gene indicates that the cell is senescent. In an embodiment, the expression of the Old-35 gene is measured by the expression of Old-35 specific RNA. In another embodiment, the expression of the Old-35 gene is measured by the expression
35 of the OLD-35 protein.

This invention also provides a method of determining whether a cell is terminally differentiated comprising measurement

-21-

of the expression of Old-35 gene, wherein the expression of the Old-35 gene indicates that the cell is terminally differentiated. In an embodiment, the expression of Old-35 gene is measured by the expression of Old-35 specific RNA.
5 In another embodiment, the expression of the Old-35 is measured by the expression of OLD-35 protein.

This invention provides a method of determining whether a cell has growth arrest comprising measurement of the
10 expression of Old-35 gene, wherein the expression of Old-35 gene indicates that the cell has growth arrest. In an embodiment, the expression of the Old-35 gene is measured by the expression of old 35 specific RNA. In another embodiment, the expression of the Old-35 gene is measured by
15 the expression of OLD-35 protein.

This invention provides a method of inhibiting growth of cancer cells comprising contacting the cancer cells with an amount of purified OLD-64 protein effective to inhibit growth
20 of cancer cells.

This invention also provides a method of inhibiting growth of cancer cells comprising contacting the cancer cells with an amount of purified OLD-64 protein effective to inhibit
25 growth of cancer cells.

This invention provides a method of determining whether a cell is senescent comprising measurement of the expression of Old-64 gene, wherein the expression of the Old-64 gene
30 indicates that the cell is senescent. In an embodiment, the expression of old-64 gene is measured by the expression of Old-64 specific RNA. In another embodiment, the expression of Old-64 gene is measured by the expression of the OLD-64 protein.

35

The expression of specific OLD RNA may be measured by the below method: (a) isolating the nucleic acids from a sample; (b) hybridizing the isolated nucleic acids with the

-22-

appropriate Old gene under conditions permitting hybrids formation; and (c) detecting the hybrid formed.

The invention provides a pharmaceutical composition for
5 inhibiting cell growth comprising a pharmaceutically acceptable carrier and purified old 35 or old 64 at a concentration effective to inhibit cell growth.

This invention provides a method of determining whether a
10 cell is terminally differentiated comprising measurement of the expression of Old-64 gene, wherein the expression of the Old-64 gene indicates that the cell is terminally differentiated. In an embodiment, the expression of Old-64 gene is measured by the expression of Old-64 specific RNA.
15 In another embodiment, the expression of Old- 64 gene is measured by the expression of the OLD-64 protein.

This invention provides a method of determining whether a cell is growth arrested comprising measurement of the
20 expression of Old-64 gene, wherein the expression of Old-64 gene indicates that the cell is growth arrested. In an embodiment, the expression of Old- 64 gene is measured by the expression of Old-64 specific RNA. In another embodiment, the expression of Old- 64 gene is measured by the expression
25 of the OLD-64 protein.

This invention provides a method of regenerating tissues comprising contacting the tissue with an inhibitor of OLD-35 or OLD-64 protein at a concentration effective to regenerate
30 said tissues.

Methods to determine such a concentration are well-known in the art. The effective concentration of said inhibitor of OLD-35 or OLD-64 protein may be determined by using different
35 concentrations of said inhibitor and examine the effect produced.

This invention provides a method of anti-aging in a cell

-23-

comprising contacting the cell with an agent for inhibiting expression of Old-35 or Old-64 gene at a concentration effective to reverse the aging process in the cell.

5 This invention provides a pharmaceutical composition for stimulating or resuming cell growth comprising a pharmaceutically acceptable carrier and purified Old-35 or Old-64 suppressant at a concentration effective to stimulate or resuming cell growth. A purified suppressant is compound
10 capable of suppressing the activity of OLD-35 or OLD-64. For example, the suppressant can act on the gene level such that no Old-35 or Old-64 gene will be switched on. Alternatively, the suppressant may be a small molecule capable of binding to the active sites on the OLD-35 or -64 protein such that
15 the protein will not be functional or the activity of the protein will decrease. A specific antibody or its binding fragment, which is capable of binding to the OLD-35 or -64, may be a suppressant.

20

This invention provides a method for screening the presence of interferon alpha or beta of a sample comprising steps of: (a) contacting the sample with cells under conditions permitting expression of Old-35 or Old-64 gene in the
25 presence of interferon alpha or beta; and (b) determining the expression of Old-35 or Old-64 gene, an increase of expression indicates the presence of interferon alpha or beta.

30 This invention provides a method for detection of the secretion of interferon alpha or beta comprising steps of: (a) obtaining an appropriate sample from the subject; and (b) detecting expression of Old-35 or Old-64 gene, the expression of Old-35 or Old-64 gene indicating the secretion of
35 interferon in a subject.

This invention provides a method for monitoring chemotherapy of a subject comprising steps of: (a) obtaining an

-24-

appropriate sample from the subject; and (b) detecting expression of Old-35 or Old-64 gene, the expression of Old-35 or Old-64 gene indicating that the chemotherapy is effective.

5 This invention provides a method for diagnosis of the proliferating stage of a tumor from a subject comprising steps of: (a) obtaining an appropriate sample from the subject; and (b) detecting expression of Old-35 or Old-64 gene, the expression of Old-35 or Old-64 gene indicating that
10 the tumor is not at a proliferating stage.

This invention also provides a kit for diagnosis of the proliferating stage of a tumor, comprising a nucleic acid molecule capable of specifically hybridizing to the nucleic
15 acid molecule of Old-35 or Old-64.

This invention also provides a kit for diagnosis of the proliferating stage of a tumor, comprising antibody capable of specifically recognizing OLD-35 or OLD-64 protein.
20

This invention provides different kits containing appropriate reagents to perform the above-described methods.

This invention also provides a method for identifying an
25 agent that modulates the expression of Old-35 or Old-64 gene, comprising: (a) contacting a candidate agent with a cell transformed or transfected with a reporter gene under the control of a Old-35 or Old-64 promoter or a regulatory element thereof under conditions and for a time sufficient
30 to allow the candidate agent to directly or indirectly alter expression of the promoter or regulatory element thereof; and (b) determining the effect of the candidate agent on the level of reporter protein produced by the cell, thereby identifying an agent that modulates expression of Old-35 or
35 64 gene.

This invention provides a method of identifying compounds that induce proliferation or cancerous phenotype,

-25-

comprising: exposing cell comprising the promoter of Old-35 or Old-64 to the compound and identifying compounds that suppress the Old-35 or 64 promoter.

5 This invention provides a method of identifying compounds that induces senescence, or terminal differentiation, comprising: exposing the cell comprising the promoter of Old-35 or Old-64 to the compound and identifying compounds that activate the Old-35 or 64 promoter.

10

This invention provides a method of identifying genes which are common to the pathway of senescence and terminal differentiation comprising steps of: (a) obtaining a subtracted library which is enriched for genes expressed in
15 terminal differentiation; (b) screening the library with senescent probe to identify novel genes which are expressed during senescence and terminal differentiation; and (c) examining the biological activity of the identified gene to determined whether it is expressed during senescence and
20 terminal differentiation.

This invention provides a method of identifying genes which are common to the pathway of senescence and terminal differentiation comprising steps of: (a) obtaining a
25 subtracted library which is enriched for genes expressed in senescence; (b) screening the library with terminal differentiation probe to identify novel genes which are expressed during senescence and terminal differentiation; and (c) examining the biological activity of the identified gene
30 to determined whether it is expressed during senescence and terminal differentiation.

This invention also provides the gene identified by the above methods.

35

This invention provides a method of degrading specific RNAs in a cell comprising induction of the expression of Old-35. This invention further provides a method of degrading

-26-

specific RNAs in a cell comprising introducing a vector into the cell comprising the Old-35 gene.

In one embodiment of the invention, expression of Old-35 can be used as diagnostic indicator of cellular senescence, terminal differentiation and/or growth suppression. Specifically, Old-35 can be used to determine if a cell has lost proliferative ability and thus has become senescent.

10 In addition, expression of Old-35 can be used as a marker to identify drugs or small molecules that will induce senescence, e.g., to inhibit cancer cell growth or abnormal proliferative states such as psoriasis, hemangioblastoma, etc..

15

Further, expression of Old-35 can be used to identify drugs or small molecules that will inhibit senescence, and thus stimulate tissue regrowth, repair and/or regeneration.

20 Still further, expression of Old-35 can be used as a marker to identify drugs or small molecules that will induce terminal cell differentiation, e.g., to inhibit cancer cell growth or abnormal proliferative states such as psoriasis, hemangioblastoma, etc..

25

Expression of Old-35 can also be used to identify drugs or small molecules that will inhibit terminal differentiation, and thus stimulate tissue regrowth, repair and/or regeneration.

30

Furthermore, expression of Old-35 can be used as marker for detecting cytokines, specifically type I interferons, in biological samples. Since type I interferons, including leukocyte and fibroblast interferons, which activate gene expression through the well characterized Jak and Stat kinase pathways, this gene (Old-35) can be used to detect or monitor drugs and other small molecules that activate these important pathways.

The combination of Old-35 with other interacting proteins can be used to target the differentiation of specific target cells, and thus result in the reprogramming of pluripotent stem cells to terminally differentiated end cells.

5

Additionally, Old-35 can be used to selectively stabilize specific mRNAs possibly containing AU rich 3' UTRs (untranslated regions). This effect can result in the sustained expression of genes potentiating or inhibiting cell growth. It could also result in the stabilizing of cytokine genes resulting in increased biological and immunological activity.

Old-35 can also be used as part of a methodology to polymerize random NTPs into nucleic acids and/or to induce the degradation of specific mRNAs.

This invention will be better understood from the Experimental Details which follow. However, one skilled in the art will readily appreciate that the specific methods and results discussed are merely illustrative of the invention as described more fully in the claims which follow thereafter.

-27-

EXPERIMENTAL DETAILS

Library Screening

A subtracted cDNA library enriched in genes modified during
5 terminal differentiation in human melanoma cells (Jiang and
Fisher, 1993) was plated at 200 pfu/plate. Colonies were
transferred to Nylon filters, denatured for 2 min (1.5M NaCl,
0.5M NaOH), neutralized for 5 min (1.5M NaCl, 0.5M Tris-HCl,
pH 8.0), and washed for 30 sec (0.2M Tris-HCl, pH 7.5, 2 X
10 SSC). Filters were cross-linked (120,000 μ J of UV energy) for
30 sec in a Strata linker (Stratagene) and prehybridized at
65°C for 2 hr in ExpressHyb (ClonTech). The probe was
denatured at 95°C for 5 min, cooled at 0°C for 5 min and then
applied to the filters at 3×10^6 cpm/ml. The filters were
15 hybridized overnight at 65°C. The next day, the filters were
washed (2 X SSC, 0.1 % SDS) 3 X for 20 min and exposed for
autoradiography.

PREPARATION OF THE PROGERIA CDNA PROBE

20 Ten μ g of total RNA derived from AG0989B cells (Progeria) (p
22) (Corriel Repository, Camden) was reverse transcribed
using SuperScript II (manufacturer's protocol, GibcoBRL)
except that 900 μ Ci of [α -³²P]-dCTP (3000Ci/mmol) (Amersham)
and 0.4mM of non-radioactive dCTP was used in place of 10mM
25 dCTP. The probe was purified using Quick Spin Columns
(Boehringer Mannheim).

PHAGE ISOLATION

The exposed film from autoradiography was aligned with the
30 phage containing plates and hybridizing clones were isolated
and re-suspended in SM buffer (1 ml).

PCR

PCR was performed for each phage stock using the
35 manufacturer's protocol (GibcoBRL) with 3 μ l of SM stock.
Since T3 and T7 primers flank the insert, these primers were
used to selectively amplify the insert from the phage vector

-28-

(Stratagene). PCR conditions were 30 cycles at 94°C for 1 min, 55°C for 1 min, 72°C for 2 min and 72°C for 10 min to allow complete extension. The PCR products were resolved on 1% agarose gels to determine the size of the product. All 5 clones were sequenced and the novel cDNAs were selected for Northern blotting analysis.

NORTHERN BLOTTING

Total RNA was extracted using the guanidinium isothiocyanate method followed by phenol/chloroform/isoamyl extraction and precipitation in isopropanol as described in Chomczynski and Sacchi (1987). The probes were labeled with [α -³²P]dCTP by random priming (Amersham). Ten μ g of total RNA were electrophoresed in a 1% agarose/2.2M formaldehyde gel and transferred to Hybond-NX filters (Amersham). Hybridization was performed in ExpressHyb solution (Manufacturer's protocol, Clontech). Briefly, filters were prehybridized at 67°C for 0.5 hr, hybridized with a denatured probe for 1.5 hr, and washed (.2 X SSC, .1 % SDS) 1 X at 24°C for 5 min, and 2 X at 55°C for 20 min.

CELLS AND CULTURE CONDITIONS

HO-1 human melanoma cells were grown in Dulbecco's modified Eagle's medium (DMEM) supplemented with 5% fetal bovine serum at 37°C in a 5% CO₂/95% air humidified incubator. Cell lines used for the senescence study were obtained from Corriel Repository (Camden, NJ). Fibroblast cell lines from patients with Progeria-Hutchinson-Gilford Syndrome (AG01972B, AG0989B, AG01178B) and normal fetal fibroblasts (GM01379A) were grown in DMEM supplemented with 15% fetal bovine serum (Gibco BRL) and 2 X essential and non-essential amino acids (Sigma). IDH4 cells (Wright et al., 1989) were grown in DMEM supplemented with 10% fetal bovine serum or 10% charcoal stripped fetal bovine serum. HO-1 cells were treated with IFN- β (2,000 U/ml) and MEZ (10 ng/ml) to induce terminal differentiation (Fisher et al., 1985). To inhibit RNA and protein synthesis, HO-1 cells were treated with actinomycin D (5 μ g/ml) and

-29-

cycloheximide (50 μ g/ml), respectively, as previously described (Jiang et al., 1993b).

STAINING FOR SENESENCE-ASSOCIATED (SA) B-GAL ACTIVITY

5 Cells were washed 2 X with PBS, fixed in 3% formaldehyde, and stained as previously described (Dimri et al., 1995). Briefly, following fixation, cells were incubated overnight at 37°C in a reaction buffer containing X-gal (1 mg/ml), 40mM citric acid/sodium phosphate (pH 6.0), potassium
10 ferrocyanide/ferricyanide (5mM), NaCl (150mM) and 2mM MgCl₂. IDH4 cells grown in the presence of dexamethasone (10⁻⁶ M) were used as a negative control.

EXPERIMENTAL RESULTS

15 Preliminary screening of cDNA libraries screening the temporally spaced subtracted differentiation inducer treated HO-1 cDNA (DISH) library enriched for genes regulated during terminal differentiation in melanoma cells, with the RNA from senescent fibroblasts, resulted in the identification of 10
20 novel and 28 known cDNAs, referred to as Old cDNAs (Table 1). Northern and reverse Northern blotting was used to determine the expression patterns of these Old cDNAs. The goal of our screening was to identify and clone differentially expressed genes common to senescence and terminal differentiation. To
25 achieve this aim, RNAs from HO-1 (untreated or treated with IFN- β , 2,000 U/ml or IFN- β (2,000 U/ml) +MEZ (10 ng/ml)), young fibroblast cultures (GM01379) and two senescent cell cultures (AG01976, AG0989B) were isolated and expression of specific Old genes was determined (Fig. 1). Since the
30 subtracted library that was screened should be enriched for HO-1 genes regulated by IFN- β and IFN- β + MEZ, it was anticipated that the level of expression of many of the Old cDNAs would be reduced or absent in actively proliferating, untreated HO-1 cells. However, since this library was
35 screened with an un-subtracted senescent probe (containing senescent specific, housekeeping and other genes) some of the cDNAs should also be expressed in non-senescent fibroblasts.

-30-

Four of the six novel cDNAs, Old-137, Old-139, Old-142 and Old-175, were expressed in both proliferating and senescent fibroblasts. Expression of two novel Old genes, Old-35 and Old-64, were restricted to the senescent fibroblasts and 5 IFN- β and IFN- β + MEZ treated H0-1 cells. Different exposure times revealed that the expression level of Old-35 is higher in senescent fibroblasts than in H0-1 cells treated with IFN- β or IFN- β + MEZ (Fig. 1). Response of Old-35 to Interferons Time-course and dose-response experiments were 10 performed in H0-1 cells to determine the temporal kinetics of Old-35 induction by IFN- β and the concentration of IFN- β capable of inducing Old-35 expression, respectively. Additionally, the effect of IFN- α , IFN- γ and TNF- α on Old-35 expression in H0-1 cells was examined. Old-35 was 15 up-regulated by IFN- β (2,000 units/ml) and IFN- β + MEZ (2,000 units/ml + 10 ng/ml) within 3 hr of treatment (Fig. 2A and D). Since IFN- β induces growth suppression in H0-1 cells at 2,000 units/ml, it was considered important to determine whether up-regulation of Old-35 could occur in the absence 20 of growth suppression. Old-35 expression was induced in H0-1 cells with as little as 1 U/ml of IFN- α , a dose of IFN that is not growth inhibitory, suggesting a direct effect of IFN on expression of this gene in the absence of growth suppression (Fig. 2B). Treatment of H0-1 cells with IFN- α 25 resulted in significant up-regulation of Old-35 in H0-1 cells, whereas this expression was marginally stimulated by IFN- γ and no detectable or consistent induction occurred with TNF- α (Fig. 2C). These experiments document differential regulation of Old-35 expression by different cytokines, with 30 type I Interferons (IFN- α /IFN- β) being the most active cytokines tested in inducing Old-35 expression in H0-1 cells. Expression of Old-35 in various human tissues and during mouse development to determine the tissue-specific expression pattern of Old-35 we examined the expression of this gene 35 using Human Multiple Tissue Northern (MTN) Blots (Clontech) (Fig. 3 A). Old-35 was expressed in all of the tissues tested, including heart, brain, placenta, lung, liver, skeletal muscle, kidney and pancreas. The highest levels of

-31-

Old-35 expression were detected in the heart and brain. Since the heart and brain contain a high proportion of non-regenerating, terminally differentiated cells, it is possible that Old-35 may be important in maintaining end stage differentiation in these target organs. Since terminal differentiation of specific tissue cell types occurs during normal development of the embryo, the expression pattern of Old-35 was determined during mouse development. The highest level of Old-35 expression was apparent during the earliest stage of development (8 days) and it steadily declined with time (10 to 16 days) (Fig. 3B). This dilution effect is frequently observed when mRNA expression is localized in a specific organ as the embryo develops, because the ratio of the region of expression to the whole body decreases over time. Since the mouse developmental Northern Blot was probed with human cDNA and the resulting signal was very strong, the homology between human and mouse OLD-35 transcripts must be very high. The EST database search showed very close homology between the mouse and the human cDNA, ~ 90% (Fig. 4).

Expression of Old-35 during growth arrest and senescence in IDH4 cell IDH4 cells were produced by transfecting IMR-90, normal human fibroblasts, with a dexamethasone (DEX) inducible mouse mammary tumor virus-driven simian virus 40 T-antigen (Wright et al., 1989). In this model system, prolonged proliferation and the absence of markers of senescence are dependent upon the continued presence of DEX and thus the SV40 T-antigen. (Wright et al., 1989). In DEX-free medium, DNA synthesis declines by ~80% within the first 3 days and reaches a minimum level at day 7. This decline corresponds with a decrease in telomerase activity and T-antigen expression (Holt et al., 1996). Since T-antigen has a long half-life (~3 days) and remains in the cells for about 5-7 days after the removal of DEX, it is possible that the up-regulation of Old-35 by day 7 corresponds with the depletion of T-antigen in these cells (Fig. 5). Further experiments to define relationship between T-antigen expression and Old-35 expression in IDH4 cells are in progress. Old-35 and p21 are coordinately expressed in

-32-

quiescent cells since many of the genes involved in terminal differentiation and senescence are predominantly active during the G₁ phase of the cell cycle, we determined whether Old-35 was expressed at this point of the cell cycle. To achieve this objective, human diploid fibroblasts were grown to confluence (a classic way to arrest and synchronize these cells) (*Tseng et al., 1983). After release of the cells from confluence, following a short lag cells re-entered G₁ phase and then the cells traversed through S, G₂, M and back to G₁. In these cells, Old-35 was highly expressed during the confluence period and at G₁ (Fig. 6). Additionally, as more of the cells entered G₁ Old-35 expression increased. After 15hr, Old-35 expression was significantly reduced, but expression increased again when the cells became confluent (20 hr). The expression of p21 (G₁ specific cyclin-dependent kinase inhibitor) coincided with the expression of Old-35 (Fig. 6).

20 STABILITY OF OLD-35 IN IFN-B TREATED CELLS

The 3' UTR of particular lymphokines, cytokines and proto-oncogenes contain ARE elements that are implicated in regulating mRNA stability (Fig. 7). The presence of four such ARE elements in the 3' UTR of Old-35 suggests that mRNA stability may contribute to differential expression of this gene under varied treatment and growth conditions. Recently, HuR a protein involved in the destabilization of mRNAs containing ARE elements has been purified and identified as a member of the Elav-line gene family (Myer et al., 1997). If the HuR protein can regulate the stability of Old-35 in HO-1 cells, then treatment of cells with cycloheximide, which inhibits protein synthesis, should decrease or eliminate the HuR protein thereby resulting in stabilization of Old-35 mRNA. Cycloheximide treatment of HO-1 cells (Fig. 8A, lane 2) and IFN-b pre-treated HO-1 cells (Fig. 8A, lane 6 and 7) increases the level of Old-35 mRNA indicating that factors responsible for its degradation might have been inhibited. The cycloheximide studies also indicate that induction of

-33-

Old-35 expression can occur in the absence of new protein synthesis (Fig. 8A, lanes 3, 4 and 5). However, since Old-35 RNA production in HO-1 cells occurs within 3 hr of treatment, and cycloheximide is present for the entire treatment period, it is possible that modifications of existing proteins may occur prior to changes in Old-35 transcription. Control of mRNA levels in a cell are regulated predominantly at two points: transcription and mRNA stability. To determine if IFN- β + MEZ or IFN- β effect Old-35 mRNA stability in HO-1 cells, the half-life of the Old-35 mRNA was determined as previously described (Jiang et al., 1993b) (Fig. 8B). Untreated and IFN- β + MEZ or IFN- β treated HO-1 cells were incubated with the RNA polymerase II inhibitor Actinomycin D (Act D) and the stability of pre-existing mRNA was determined by Northern blotting. This experiment revealed that the half-life of Old-35 mRNA in HO-1 cells treated with IFN- β + MEZ or IFN- β is ~6-8 hr, suggesting that stabilization of this mRNA may contribute to the elevation of Old-35 levels in treated cells (Fig. 8B and data not shown). However, because of the low level of Old-35 expression in untreated HO-1 cells, it was not possible to accurately determine the half-life of this message in these cells. Whether the observed low levels of Old-35 mRNA in untreated actively proliferating HO-1 cells are the result of a lack of transcriptional activation or mRNA stability still remain to be determined. Nuclear run-on assays, that measure rate of RNA transcription, should reveal whether the promoter is active in HO-1 cells in the absence of IFN- β treatment and the potential contribution of transcriptional activation to elevated Old-35 mRNA following IFN- β and IFN- β + MEZ treatment.

CLONING AND SEQUENCE ANALYSIS OF OLD-35

An initial 600bp fragment of Old-35 was identified and cloned from a differentiation inducer treated subtracted (DISH) HO-1 cDNA library as described in the library screening protocol. This cDNA was cloned in a pBlueScript vector in the opposite orientation 3'-5' (EcoRI-XhoI) as a result of subtraction

-34-

hybridization. During the subtraction procedure, cDNAs are excised from the vector by double-digestion with EcoRI and XhoI. Since many cDNA also contain internal EcoRI-XhoI sites, many cDNAs will be cut internally and after the subtraction 5 procedure they will re-ligate in the incorrect direction. Thus the original 600bp fragment of Old-35 contained an internal region of Old-35 cDNA and lacked 3' and 5' flanking sequences. The 5' region of Old-35 was cloned from IFN- β treated HO-1 cells using a recently developed cDNA extension 10 procedure, complete open reading frame cloning (C-ORF), yielding in a single-reaction an ~2kb fragment (Kang and Fisher, unpublished). The 3' region of Old-35 was cloned using the 3' RACE procedure with 3' gene specific nested primers and dT, yielding an ~400bp product. The final 15 sequence of Old-35 is shown in Fig. 9. Although a portion of the 5' may still be missing, the Old-35 cDNA obtained using C-ORF and 3' RACE represents a near full-length clone judging from the Northern blotting data (Fig. 1), in which Old-35 hybridizes with an ~2.4-2.7 kb RNA species. Sequence 20 analysis revealed that the Old-35 cDNA (~2.6kb) contains a less frequently observed polyadenylation site (AUUAAA) (found in only ~10% of cDNAs) (Manley et al., 1988). The putative protein sequence does not exhibit homology to any known genes except to the *Escherichia coli* PNPase (polyribonucleotide 25 phosphorylase) gene of which 30% of the sequence is homologous and 50% displays sequence similarity (Fig. 10).

EXPERIMENTAL DISCUSSION

Controlled cellular proliferation is paramount for sustaining 30 homeostasis in multicellular organisms. The regulation of this dynamic process is of particular relevance in maintaining a balance between cell loss and cell renewal, important factors in development, differentiation and aging. Moreover, abnormalities in cell division are hallmarks of 35 many disease states, including developmental and congenital birth defects, premature aging syndromes and abnormal proliferative states such as cancer. Several genes involved

-35-

in cell proliferation control, including the tumor suppressor p53 and the cyclin dependent kinase (cdk) inhibitor p21, display elevated expression in growth suppressive conditions, such as quiescence (Niculescu et al., 1998, Lacombe et al., 5 1996, Linke et al., 1996), senescence (Irving et al., 1992; Gire and Wynford-Thomas, 1998) and terminal cell differentiation (Jiang et al., 1994b, 1995b; Steinman et al., 1994). Since both terminal differentiation and senescence are characterized by growth arrest, it is possible that similar 10 and overlapping genes and gene expression changes may mediate these processes. To test this hypothesis we have screened a subtracted differentiation inducer treated human H0-1 melanoma library with mRNA derived from senescent human fibroblasts. This approach has resulted in the isolation of 15 a large number of cDNAs, consisting of both known and novel sequences (Table 1), displaying elevated expression in senescent human fibroblasts. Several of the same cDNAs, have also been independently identified from the same subtracted H0-1 library after screening with mRNA isolated from H0-1 20 cells treated with IFN- β + MEZ that induce irreversible growth arrest and terminal differentiation (Huang et al., 1999). This observation validates our hypothesis and suggests that this novel approach may prove useful in identifying and cloning genes displaying coordinated expression as a function 25 of induction of growth arrest during terminal differentiation and cellular senescence. One such cDNA is the novel gene, Old-35.

Induction of terminal differentiation in human melanoma cells 30 by IFN- β + MEZ frequently results in the induction and up-regulation of genes that also display elevated expression following exposure to IFN- β , referred to as Type I melanoma differentiation associated (*mda*) genes (Jiang and Fisher, 1993). Old-35 represents such a gene, since its expression 35 is elevated in H0-1 cells after treatment with IFN- β and IFN- β + MEZ. Old-35 is also up-regulated during growth arrest and senescence in human fibroblasts, indicating that its expression is not restricted to only programs of

-36-

differentiation or to human melanoma cells. Since IFN- β has well-established antiproliferative properties, in both normal and cancer cells (Fisher and Grant, 1985), it is possible that Old-35 may function as a down-stream gene in the 5 IFN-signaling pathway culminating in growth arrest. A number of experiments indicate that Old-35 expression is related to cellular senescence and proliferative quiescence. Analysis of Northern blots from young versus senescent human fibroblasts indicates restricted expression of Old-35 to 10 senescent cells. IDH4 cells, conditionally immortalized by a DEX-inducible SV40 T-antigen, represent an excellent in vitro model to study senescence (Wright et al., 1989). The presence of DEX in the growth media allows the IDH4 cells to actively proliferate, while the absence of it causes them to 15 senesce. In these cells, Old-35 expression is only detected after 7 days of growth in media devoid of DEX. This expression also corresponds with the SA- β (-GAL staining of IDH4 cells, a well-established senescence marker (Dimri et al., 1995). Old-35 expression also increases when fibroblasts 20 become arrested in G₀ by growth and maintenance at confluence. In these contexts, Old-35 could prove useful as a diagnostic marker for cellular senescence, terminal differentiation and growth arrest. High levels of Old-35 expression are also found in the brain and heart, the only 25 human tissues that do not possess active regenerative properties. Judging from the localized expression of Old-35 during development, this gene may contribute to heart and brain development by assisting in the maintenance of terminal differentiation of cells in these organs. Due to the high 30 sequence homology of Old-35 to bacterial polyribonucleotide phosphorylase (PNPase), it is possible that Old-35 protein may exhibit a PNPase enzymatic activity. PNPase is one of the critical components of the *Escherichia coli* RNA degradosome (Blum et al., 1997), which consists of both PNPase and 35 endoribonuclease RNase E. The function of this complex is to control the rate of mRNA degradation. The PNPase possesses two enzymatic activities, 3'-5' processive exoribonuclease

-37-

activity and 5'-3' RNA polymerase activity (Blum et al., 1997). Recently, it has been shown that PNPase also has the capacity to bind to a specific double-stranded DNA sequence in a sequence-specific manner (Zhang et al., 1998). Since
5 Old-35 is differentially expressed in cells that undergo growth arrest, it is possible that this gene may play a role in RNA degradation in growth arrested cells. Additionally, since genes containing AUUUA elements (Myer et al., 1997) have been shown to be involved in the global regulation of
10 gene expression, it is possible that Old-35 by binding sequence-specific targets, controls growth related gene expression. In this context, Old-35 might display tumor suppressor properties and could be useful for the gene therapy of cancer.

-38-

TABLE 1

CLONE DESIGNATION	CLONE IDENTITY
old-1	Vimentin
5	old-2 Human ribosomal protein S3a, v-fos
old-5	mRNA M phase phosphoprotein
10	old-7 RIG-G, Cig49
old-11	MHC class I lymphocyte antigen
old-14	Human non-muscle myosin alkaline light chain
15	old-18 Human ADP-ribosylation factor 4
old-19	Human mitochondrial cytochrome oxidation
20	old-24 56 kDa IFN inducible
old-30	Ribosomal protein L5
old-32*	Novel*
25	old-34 IFN-inducible protein
old-35*	Novel*
30	old-38 H.s. small acidic protein
old-39	Human acidic ribosomal phosphatase
old-42	Neurofibromatosis type 1

-39-

	Old-59	Human nuclear receptor hTAK1
	Old-60	Mitochondrial DNA
5	Old-61	Transcription factor I (99%)
	Old-64*	Novel*
10	Old-65	CDC16HS cell 81, 261-68
	Old-74	Human ISG 54K gene (IFN-gamma)-cig42
	Old-79	Human T-complex polypeptide I gene
15	Old-80	Vitamin D induced
	Old-83*	Novel*
	Old-87*	Novel*, Possibly similar to Old-83
20	Old-107*	Novel*-Human homologue of Cow G-Protein
	Old-113	DNA binding protein
25	Old-115	U1 small nuclear RNP
	Old-119	Human HS1 protein
30	Old-121*	Novel*
	Old-137*	Novel*
	Old-139*	Novel*

-40-

5

10

Old-140	Human putative trans. CA150
Old-142*	Novel*
Old-144	MLN70 calcium- binding
Old-165	T-cell cyclophilin
Old-170	Human homologue of rat zinc transporter
Old-175 (5-3)*	Novel*

-41-

REFERENCES

- Blum E., Py B., Carpousis A.J., and Higgins C.F. (1997). Polyphosphate kinase is a component of the Escherichia coli RNA degradosome. Mol. Microbiol., 26: 387-398.
- 5
- Campisi, J., (1996). Replicative Senescence An Old Lives Tale Cell. Cell 84(4):497-500.
- Caput D., Beutler B., Hartog K., Thayer R., Brown-Shimer S.,
10 and Cerami A. (1986). Identification of a common nucleotide sequence in the 3'-untranslated region of mRNA molecules specifying inflammatory mediators. Proc Natl Acad Sci U S A, 83: 1670-1674.
- 15 Chen C.Y., and Shyu A.B. (1995). AU-rich elements: characterization and importance in mRNA degradation. Trends Biochem Sci., 20: 465-470. Chomczynski P., and Sacchi N. (1987). Single step method of RNA isolation by acid guanidinium thiocyanate-phenol-chlorophorm extraction. Anal.
20 Biochem., 162: 156-159.
- Darnell J.E. Jr., Kerr I.M., and Stark G.R. (1994). Jak-STAT pathways and transcriptional activation in response to IFNs and other extracellular signaling proteins. Science, 264:
25 1415-1421.
- Dimri G.P., Lee X., Basile G., Acosta M., Scott G., Roskelley C., Medrano E.E., Linskens M., Rubelj I., Pereira-Smith O., Peacocke M., and Campisi, J. (1995). A biomarker that
30 identifies senescent human cells in culture and in aging skin in vivo. Proc. Natl. Acad. Sci. USA, 92: 9363-9367.
- Fisher P.B., and Grant S. (1985). Effects of interferon on differentiation in normal and tumor cells. Pharmacology and
35 Therapeutics, 27: 143-166. Fisher P.B., Prignoli D.R., Herms H. Jr., Weinstein I.B., and Pestka S. (1985). Effects of

-42-

combined treatment with interferon and mezerein on melanogenesis and growth in human melanoma cells. J. Interferon Res., 5: 11-22.

5 Gire V., and Wynford-Thomas D. (1998). Reinitiation of DNA synthesis and cell division in senescent human fibroblasts by microinjection of anti-p53 antibodies. Mol. Cell. Biol., 18: 1611-1621.

10 Goeddel D.V., Leung D.W., Dull T.J., Gross M., Lawn R.M., McCandliss R., Seeburg P.H.,

Goldstein, S., (1990) Replicative Senescence The Human Fibroblast Comes Of Age. Science 249:1129-1133.

15

Gutterman J.U. (1994). Cytokine therapeutics: lessons from interferon alpha. Proc. Natl. Acad. Sci. USA, 91: 1198-1205.

Hayflick L., and Moorehead P.S. (1961). The serial
20 cultivation of human diploid cell strains. Exp. Cell Res., 25: 585-621.

Holt S.E., Wright W.E., and Shay J.W. (1996). Regulation of telomerase activity in immortal cell lines. Mol. Cell. Biol.,
25 16: 2932-2939.

Huang F., Adelman J., Jiang H., Goldstein N.I., and Fisher P.B. (1999). Identification and temporal expression pattern of genes modulated during irreversible growth arrest and
30 terminal differentiation in human melanoma cells. Oncogene, in press.

Irving J., Feng J., Wistrom C., Pikaart M., and Villeponteau B. (1992). An altered repertoire of fos/jun
35 (AP-1) at the onset of replicative senescence. Exp Cell Res., 202: 161-166.

-43-

Jiang H., and Fisher P.B. (1993). Use of a sensitive and efficient subtraction hybridization protocol for the identification of genes differentially regulated during the induction of differentiation in human melanoma cells. Mol. Cell. Different., 1: 285-299.

Jiang H., Lin J., and Fisher, P. B. (1994a). A molecular definition of terminal cell differentiation in human melanoma cells. Mol. Cell. Different., 2: 221-239.

10

Jiang H., Lin J., Su Z.-z., Collart F.R., Huberman E., and Fisher P.B. (1994b). Induction of differentiation in human promyelocytic HL-60 leukemia cells activates p21, WAF1/CIP1, expression in the absence of p53. Oncogene, 9: 3397-3406.

15

Jiang H., Lin J.J., Su Z.-z., Goldstein N.I., and Fisher P.B. (1995a). Subtraction hybridization identifies a novel melanoma differentiation associated gene, mda-7, modulated during human melanoma differentiation, growth and progression. Oncogene, 11: 2477-2486.

20

Jiang H., Lin J., Su Z.-z., Kerbel R.S., Herlyn M., Weissman R.B., Welch D., and Fisher P.B. (1995b). The melanoma differentiation associated gene mda-6, which encodes the cyclin-dependent kinase inhibitor p21, is differentially expressed during growth, differentiation and progression in human melanoma cells. Oncogene, 10: 1855-1864.

25

Jiang H., Su Z.-z., Boyd J., and Fisher P.B. (1993a). Gene expression changes associated with reversible growth suppression and the induction of terminal differentiation in human melanoma cells. Mol. Cell. Different., 1: 41-66.

30

Jiang H., Waxman S., and Fisher P.B. (1993b). Regulation of c-fos, c-jun and jun-B gene expression in human melanoma cells induced to terminally differentiate. Mol. Cell. Different., 1: 197-214.

35

-44-

- Kashima N., Nishi-Takaoka C., Fujita T., Taki S., Yamada G., Hamuro J., and Taniguchi T. (1985). Unique structure of murine interleukin-2 as deduced from cloned cDNAs. Nature, 313: 402-404.
- 5 Lacombe L., Orlow I., Silver D., Gerald W.L., Fair W.R., Reuter V.E., and Cordon-Cardo C. (1996). Analysis of p21WAF1/CIP1 in primary bladder tumors. Oncol. Res., 8: 409-414.
- 10 Lee, W.M. (1988) Linc, Currant Activation Of The Transforming Potential Of The Human Fos Proto-Oncogene Requires Message Stabilization And Results In Increased Amounts Of Partially Modified Fos Protein. Mol. Cell. Biol. 8(12)5521-5527.
- 15 Lin J.J., Jiang H., and Fisher P.B. (1996). Characterization of a novel melanoma differentiation associated gene, mda-9, that is down-regulated during terminal cell differentiation. Mol. Cell. Different., 4 (4): 317-333.
- 20 Lin J.J., Jiang H. and Fisher P.B. (1998). Melanoma differentiation associated gene-9 is a human gamma interferon responsive gene. Gene, 207: 105-110.
- 25 Linke S.P., Clarkin K.C., Di Leonardo A., Tsou A., and Wahl G.M. (1996). A reversible, p53-dependent G₀/G₁ cell cycle arrest induced by ribonucleotide depletion in the absence of detectable DNA damage. Genes & Devel., 10(8): 934-947.
- 30 Murano S., Thweatt R., et al., (1991) Diverse Gene Sequences Are Overexpres. In Werner Syndrome Fibroblasts Undergoing Premature Replicative Senescence. Mol. Cell. Biol. 11(8):39005-39014.
- 35 Myer V.E., Fan X.C., and Steitz J.A. (1997). Identification of HuR as a protein implicated in AUUUA-mediated mRNA decay.

-45-

EMBO J., 16: 2130-2139.

Medrano E.E., Yang F., Boissy R., Farooqui J., Shah V., Matsumoto K., Nordlund J.J., and Park H.Y. (1994). Terminal 5 differentiation and senescence in the human melanocyte: repression of tyrosine-phosphorylation of the extracellular signal-regulated kinase 2 selectively defines the two phenotypes. Mol. Biol. Cell., 5: 497-509.

10 Nedwin G.E., Naylor S.L., Sakaguchi A.Y., Smith D., Jarrett-Nedwin J., Pennica D., Goeddel D.V., and Gray P.W. (1985). Human lymphotoxin and tumor necrosis factor genes: structure, homology and chromosomal localization. Nucleic Acids Res., 13:6361-6373.

15

Niculescu A.B., Chen X., Smeets M., Hengst L., Prives C., and Reed S.I. (1998). Effects of p21(Cip1/Waf1) at both the G₁/S and the G₂/M cell cycle transitions: pRb is a critical determinant in blocking DNA replication and in preventing 20 endoreduplication. Mol. Cell. Biol., 18: 629-643.

Sambrook, J., et al.(1989) Molecular Cloning: a laboratory manual. Second Edition. Cold Spring Harbor Laboratory Press (1989).

25

Shaw G., and Kamen R. (1986). A conserved AU sequence from the 3' untranslated region of GM-CSF mRNA mediates selective mRNA degradation. Cell, 46: 659-667.

30 Smith J.R., and Pereira-Smith O.M. (1996) Replicative senescence: implications for in vivo aging and tumor suppression. Science, 273: 63-67.

Stark G.R., Kerr I.M, Williams B.R.G., Silverman R.H., and 35 Schreiber, R.D. (1998). How cells respond to interferons. Annu. Rev. Biochem., 67: 227-264

-46-

- Steinman R.A., Hoffman B., Iro A., Guillouf C., Liebermann D.A., and el-Housein ME. (1994). Induction of p21 (WAF-1/CIP1) during differentiation. Oncogene, 9: 3389-3396.
- 5 Ullrich A., Yelverton E., and Gray P.W. (1981). The structure of eight distinct cloned human leukocyte interferon cDNAs. Nature, 290: 20-26.
- Van Straaten F., Muller R., Curran T., Van Beveren C., and
10 Verma I.M. (1983). Complete nucleotide sequence of a human c-onc gene: deduced amino acid sequence of the human c-fos protein. Proc Natl Acad Sci. USA, 80: 3183-3187.
- Wong G.G., Witek J.S., Temple P.A., Wilkens K.M., Leary
15 A.C., Luxenberg D.P., Jones S.S., Brown E.L., Kay R.M., Orr E.C., Shemaker C., Golde D.W., Kaufman R.J., Hewick R.M., Wang E.A., and Clark S.C. (1985). Human GM-CSF: molecular cloning of the complementary DNA and purification of the natural and recombinant proteins. Science, 228: 810-815.
- 20 Wright W.E., Pereira-Smith O.M., and Shay, J.W. (1989). Reversible cellular senescence: implications for immortalization of normal human diploid fibroblasts. Mol. Cell. Biol., 9: 3088-3092.
- 25 Zhang P., Vigne J.L., and Mellon S.H. (1998). Polyribonucleotide phosphorylase is a double-stranded DNA-binding protein. DNA Cell Biol., 17: 169-175.

-47-

Example #1**Background and significance**

During terminal differentiation and senescence many genes are differentially expressed. Two processes that control the overall mRNA levels are transcription and mRNA stability. Since both proliferation and differentiation are dynamic processes requiring continuous regulation (Blau, H.M., 1992, Blau et al., 1992, Blau et al, 1985) a thorough knowledge of the molecular mechanisms that regulate gene expression will significantly contribute to our understanding of development, differentiation and malignancy. Gene expression is regulated by two mechanisms: transcriptional mechanisms which determine the rate of mRNA production and equally important but under-studied post-transcriptional mechanisms which determine the overall amount of protein being produced. The experimental data from *Xenopus laevis*, *Drosophila melanogaster*, *Caenorhabditis elegans* document the importance of post-transcriptional mechanisms in early patterning of the embryos which directs correct distribution, stability, and translation of inherited maternal transcripts (Seydoux, G., 1996) Additionally, in plants, it has been shown that it is the post-transcriptional regulation and not transcription that directs the differentiation of chloroplast from its protoplast precursor (Deng and Gruissem, 1987). In mammals, posttranscriptional regulation appears to be important in cells responding to environmental stress, proliferation and differentiation (June et al, 1990, Sierra et. al., 1994)

The sequences responsible for post-transcriptional regulation are found in the 3' untranslated regions (3'UTR) of the transcripts. When orthologous genes were compared, large regions were found to exhibit more than 70% conservation over 300-500 million years of evolution, from mammals, birds, amphibians, or fish (Spicher et al., 1998).

-48-

Post-transcriptional regulation of mRNA levels is a pivotal control point in gene expression. Early response genes, such as cytokines, lymphokines and proto-oncogenes are regulated by a cis-acting adenylate-uridylate-rich element 5 (ARE) found in the 3' untranslated region (UTR) of the mRNA (Caput et al., 1986; Shaw and Kamen, 1988; Chen and Shyu, 1995; Myer et al., 1997). Currently, three classes of destabilizing elements have been identified: AUUUA-lacking elements and AUUUA-containing elements grouped into those 10 with scattered AUUUA motifs (such as proto-oncogenes) and those with overlapping AUUUA motifs (such as growth factors) (Chen et al., 1995; Myer et al., 1997). A replacement of 3'UTR containing ARE in place of a 3'UTR of a stable message, such as β -globin or luciferase targets 15 this very stable mRNA for rapid degradation (Shaw and Kamen, 1988, Maddireddi et al., 2000). In contrast, the removal of an ARE stabilizes an otherwise labile message (Miller et al., 1984; Lee et al., 1988)

A pool of genes involved in mRNA stability remains very 20 small. However, one of the best studied family of genes in this area is Elav. Elav, which stands for embryonic -lethal abnormal vision, was first identified in *Drosophila melanogaster*. Deletion mutants of the elav gene are embryonic lethal because of lack of neuronal 25 differentiation (Robinow and White, 1991). In mammals and in *Xenopus*, the elav gene family consists of three members that are developmentally regulated and tissue specific (Hel-N1, HuC, HuD,) and one member that is ubiquitously expressed called HuR (Szabo et al, 1991, Good, 1995, Ma et 30 al., 1996, Antic and Keene, 1997). The mechanism by which Elav genes promote the differentiation of neurons is not completely understood, however, it is known that Elav can bind AU rich elements in ther 3'UTRs of selected genes. By selectively stabilizing selected genes, the overall amount 35 of gene expression changes observed during terminal differentiation of neurons is regulated.

-49-

To obtain further insights into 3' UTR stabilization, 3' end maturation has been studied in detail in plants and bacteria. It is worth noting that the protein complexes involved in this process in these two completely different organisms are highly conserved. They are composed of endonucleases, exonucleases, helicases and enolases. *E.coli*, which lives in an energy high environment, has two exonucleases involved in the processing of 3'UTRs: RNase II, which has hydrolytic activity and PNPase (polynucleotide phosphorylase) which has phosphorolytic activity (Higgins et al., 1993). Single mutant of either PNPase or RNase II is viable, whereas double mutants die (Donovan and Kushner, 1986). On the other hand, *B.Subtilis*, which lives in the soil-an energy poor environment- exclusively uses PNPase and lacks RNase II. There may be a few different explanations for the presence of two exonucleases in *E.coli*. Firstly, the two exonucleases may have different specificities. This is supported by the fact that a specific degradation of S20 mRNA accumulates in pnp mutants but it fails to accumulate in rnb (Rnase II) mutants (Mackie, 1989). Another explanation could be that PNPase is phosphorolytic and Rnase II is hydrolytic. As phosphorolysis releases nucleotide diphosphates (Gedefroy-Colburn and Grunberg-Manago, 1966), the energy of the phosphate bond is conserved. Differential use of these two enzymes may reflect an adaptation to changing energy conditions (Deutscher and Reuven, 1991). This model is supported by the fact that *B. subtilis*, which normally inhabits low energy environment, uses PNPase exclusively, while *E.coli* predominantly uses Rnase II. Another interesting point worth noting is that PNPase also functions during competence development of *B.subtilis*. Since competence is a state during which specialization is acquired, competence has been used as a simple model for differentiation. Genetic competence may be defined as a physiological state enabling a bacterial culture to bind and take up high-molecular-weight exogenous DNA

-50-

(transformation). The study of competence genes has permitted their classification into two broad categories. Late competence genes are expressed under competence control and specify products required for the binding, uptake, and processing of transforming DNA. Regulatory genes specify products that are needed for the expression of the late genes (Dubnau, 1991). PNPase is necessary for the expression of late competence genes. Transformability of pnp mutant is 1-5% of that seen in wild type strains (Luttinger A, et al., 1996)

In plants, PNPase functions during chloroplast differentiation where it is involved in processing of plastid 3' UTR (example: petD). It is interesting to note that plastid genes also possess AU rich regions in its 3' UTR. Identically to bacteria, plant PNPase has a 3'-5' processive exonuclease activity that exhibits increased specificity for poly(A) and poly(U). (Hayes et al., 1996)

Human teratocarcinoma cells (NT2) can be differentiated into neurons with retinoic acid treatment and thus provides an excellent model to study neuronal differentiation. Recently it has been shown that a member of elav family, Hel-N1, when transfected into NT2 cells, forms neurites, an early sign of differentiation. However it does not cause terminal differentiation (Antic et al., 1999). Since Old-35 encodes PNPase, a 3'-5' exonuclease involved in degradation of mRNA sequences, it is possible that Old-35 can increase the effects of Hel-N1 in NT2 cells and cause them to differentiate.

30

Determining the half-life of Old-35 mRNA in HO-1 cells

Since Old-35 has an AU rich 3'UTR (Fig.4) we have speculated that its expression may be regulated by post-transcriptional mechanisms. One way to study post-transcriptional processes is to investigate mRNA half-lives. In a mammalian cell culture system this can be achieved by treating cells with Actinomycin D (AD). Since

-51-

AD inhibits RNA polymerase II activity, mRNA synthesis is terminated and the mRNA synthesized before AD treatment is allowed to decay. Total RNA is collected at different time points and quantified using Northern analyses. Using this 5 protocol, we have examined the half-lives of Old-35 mRNA in HO-1, confluent HO-1, IFN- β treated, and IFN- β +MEZ treated HO-1. The half-life of Old-35 in all the treatments did not change and (Figure 11) was estimated to be 6 hr. Since there was no difference in half-life between HO-1 and IFN- β 10 treated HO-1 it is assumed that a post-transcriptional mechanism is not responsible for the upregulation of Old-35 mRNA level in IFN- β treated HO-1.

Expression of Old-35 during growth arrest and senescence of 15 IDH4 and AR5 cells

IDH4 cells were produced by transfecting IMR-90, normal human fibroblasts, with a dexamethasone (DEX) inducible mouse mammary tumor virus-driven simian virus 40 T-antigen (Wright et al., 1989). In this model system, 20 prolonged proliferation and the absence of markers of senescence are dependent upon the continued presence of DEX and thus the SV40 T-antigen. (Wright et al., 1989). In DEX-free medium, DNA synthesis declines by ~80% within the first 3 days and reaches a minimum level at day 7. This 25 decline corresponds with a decrease in telomerase activity and T-antigen expression (Holt et al., 1996). Since T-antigen has a long half-life (~3 days) and remains in the cells for about 5-7 days after the removal of DEX, it is possible that the up-regulation of Old-35 by day 7 30 corresponds with the depletion of T-antigen in these cells (Figure 12). However, there is one drawback associated with using IDH4 cells. Since the expression of T-antigen is dependent upon DEX, a shift of IDH4 cells towards senescence is dependent upon a complete depletion of DEX 35 from the media and serum in which the cells are growing. This is normally achieved by charcoal stripping of the

-52-

serum. However, since fetal serum contains vast amount of steroids, it becomes a challenge to do so in completion. Thus, the reproducibility of complete DEX depletion is a problem. To overcome this problem, we used another cell line, AR5. AR5 is very similar to IDH4, except the fact that T-antigen is not DEX inducible but it is rather temperature sensitive. AR5 cells are able to grow rapidly at 35°C since they are expressing T-antigen. When shifted to 39°C, T-antigen is degraded and the cells become senescent. Total RNA was collected from AR5 cells grown at 35°C and from AR5 cells shifted to 39°C. Old-35 was expressed one day after the shift and at the later time points as well (**Figure 12**). To make sure that the cells had reached senescence when shifted to 35°C, we hybridized the Northern blot to the well characterized senescence marker, p21 (CDK inhibitor) (**Figure 12**). p21 expression increased in AR5 cells shifted to 39°C and showed a pattern similar to Old-35.

The difference between expression of Old-35 in IDH4 and AR5 cells can be accounted for by the differences in T-antigen depletion. Since T-antigen degrades much faster in AR5 cells (temperature sensitive) than in IDH4 cells (half-life 2-3 days), AR5 cells reach a senescent state at much faster rate than DEX depleted IDH4 cells.

25

Cloning of the second variant (3.8 kb)

Once most of the sequence was known, the cDNA was screened against the BLAST-EST database. In this search we have identified another version of Old-35 (3.8 kb) which is probably the upper band observed on Northern blots. The 3.8 kb EST was sequenced. The sequence analysis revealed differences in 3' UTRs of the 2.6kb and 3.8kb fragments. This may result from different polyadenylation patterns. To make sure that the upper band on the Old-35 Northern Blot represents the 3.8kb fragment, we will use the 3' UTR of the ATCC clone as a probe (**Figure 13**)

-53-

Old-35-GFP localization

Since no antibody for Old-35 is currently available, we decided to test the localization of Old-35 by creating an N-terminal fusion of Old-35 and GFP (Clontech). Old-35
5 was cloned in frame with GFP without the first ATG and then transfected into HeLa and HO-1 cells with SuperFect reagent (Clontech). The protein was allowed to express for 24hr. As expected for a degradative enzyme, Old-35 localized to the cytoplasm of HeLa (Figure 14) and HO-1 (data not shown).

10

Expression of Old-35 during mouse development

Using Human Multiple Tissue Northern (MTN) Blots (Clontech) we determined that Old-35 was expressed in all of the tissues tested with the highest levels in the heart
15 and brain. Since terminal differentiation of specific tissue cell types occurs during normal development of the embryo, the expression pattern of Old-35 was determined during mouse development. The highest level of Old-35 expression was apparent during the earliest stage of
20 development (8 days) and it steadily declined with time (10 to 16 days). To determine spacial expression of Old-35, in situ hybridization experiments were performed. Murine Old-35 was expressed in the spinal tube and in the arteries. However more experiments have to be performed to correctly
25 determine the expression pattern (Figure 15)

Effect of different interferon- α subtypes on Old-35 expression of Old-35

All subtypes of IFN- α stimulated Old-35 expression.
30 IFN- α H and α I-stimulated Old-35 in the lowest extent. (Figure 16). The above experiments document differential regulation of Old-35 expression by different cytokines, with type I interferons (IFN- α /IFN- β) being the most active cytokines tested in inducing Old-35 expression in
35 HO-1 cells. Moreover, since IFN signaling cascades

-54-

include Jak and Stat activation, they may prove to be important intermediates of Old-35 induction and expression.

5 Old-35 genomic structure

As described above, we have identified two BACs that were 100% homologous to specific regions of Old-35 cDNA. First BAC (RPCI-11, Plate=702,Col=8, Row=C) (Research Genetics) showed 100% homology to the 2207-2365 region of Old-35 cDNA. The sequenced regions flanking the Old-35 sequence were foreign thus it is highly possible that they are introns. The second BAC (CITBI-E1, clone 2505G20) (Research Genetics) showed 100% homology in 235bp-313bp region of Old-35 cDNA. After sequencing of the BACs, it became apparent that the Old-35 gene is distributed among 28 exons (Table 2). The spaces in the intron column signify no data for the intron size. The intron sizes are being determined.

Interestingly, there are at least three pseudogenes of Old-35 in the human genome. The first one is 92% homologous to the Old-35 cDNA and contains a portion of the cDNA (48bp-1387bp). 5' and 3' ends of the cDNA could not be found on this BAC. The second pseudogene is present on the 3rd chromosome as determined by BLAST search at it contains a cDNA fragment from the 49th nucleotide to the end of cDNA. This pseudogene exhibits 92% homology to the Old-35 cDNA. The third pseudogene also contains a cDNA fragment from 49 bp to 2517 bp. The second and third BACs are 90% homologous. In all cases, all of the BACs are highly mutated and intronless parts of the Old-35 cDNA.

TABLE 2

EXON-INTRON STRUCTURE

OF OLD-35

exons	exon size	introns	intron size	exons	exon size	introns	intron size
1	174	1	6000	15	36	15	
2	60	2	1100	16	66	16	
3	74	3	1300	17	89	17	
4	105	4	1100	18	53	18	
5	49	5		19	105	19	
6	63	6		20	72	20	
7	47	7		21	63	21	
8	113	8	6600	22	83	22	
9	186	9	800	23	83	23	
10	51	10	600	24	106	24	
11	57	11	3500	25	55	25	
12	96	12	800	26	77	26	
13	102	13		27	45	27	
14	70	14		28	406		

-56-

What is claimed is:

1. An isolated nucleic acid molecule encoding an OLD-35 or OLD-64 protein.
- 5 2. The isolated nucleic acid molecule of claim 1 wherein the nucleic acid comprises a nucleic acid having a sequence substantially the same as set forth in SEQ. ID. No.39 or 19.
- 10 3. An isolated nucleic acid molecule encoding an OLD-137, OLD-139, OLD-142, or OLD-175 protein.
4. The isolated nucleic acid molecule of claim 3 wherein the nucleic acid comprises a nucleic acid having a
15 sequence substantially the same as set forth in SEQ. ID. Nos.31, 32, 34 or 38.
5. An isolated nucleic acid molecule of claim 1, or 3, wherein the nucleic acid is DNA, genomic DNA, cDNA,
20 synthetic DNA, or RNA.
6. A nucleic acid molecule comprising a nucleic acid molecule of at least 15 nucleotides capable of specifically hybridizing with a sequence included
25 within the sequence of the nucleic acid molecule of claim 1, or 3.
7. A nucleic acid molecule of claim 6 wherein the nucleic acid is DNA, genomic DNA, cDNA, synthetic DNA or RNA.
30
8. An antisense nucleic acid molecule comprising a sequence complementary to the nucleic acid of claim 1 or 3.
- 35 9. The antisense nucleic acid molecule of claim 8, capable of inhibiting the expression of the hybridized

-57-

gene.

10. An isolated nucleic acid molecule of claim 1, 3, or 8
operatively linked to a promoter of RNA transcription.
- 5 11. A vector which comprises the isolated nucleic acid
molecule of claim 1, 3 or 8.
12. A host vector system for the production of a protein
10 having the biological activity of OLD-35 or OLD-64
protein which comprises the vector of claim 11 in a
suitable host.
13. A host vector system for the production of a protein
15 having the biological activity of OLD-137, OLD-139,
OLD-142, OLD-175 protein which comprises the vector of
claim 11 in a suitable host.
14. A method of producing a protein having the biological
20 activity of OLD-35, OLD-64 OLD-137, OLD-139, OLD-142,
OLD-175 protein which comprises growing the host
vector system of claim 12, or 13 under conditions
permitting production of the protein and recovering
the protein so produced.
- 25 15. A purified, OLD-35 protein.
16. A purified, OLD-64 protein.
- 30 17. A purified, OLD-137 protein.
18. A purified, OLD-139 protein.
19. A purified, OLD-142 protein.

35

-58-

20. A purified, OLD-175 protein.
21. A protein encoded by the isolated nucleic acid molecule of claim 1 or 3.
- 5 22. An antibody or antigen-binding fragment thereof that specifically binds to OLD-35, OLD-64, OLD-137, OLD-139, OLD-142 or OLD-175 protein.
- 10 23. A monoclonal antibody of claim 22.
24. A method of inhibiting growth of cancer cells comprising contacting the cancer cells with an amount of purified OLD-35, or OLD-64 protein or a portion thereof effective to inhibit growth of cancer cells.
- 15 25. A method for reversing the cancerous phenotype of a cancer cell which comprises introducing a nucleic acid comprising an Old-35 or Old-64 gene or a portion thereof into the cell under conditions permitting the expression of the gene so as to thereby reverse the cancerous phenotype of the cell.
- 20 26. A method for reversing the cancerous phenotype of a cancer cell in a subject which comprises introducing a nucleic acid molecule comprising an Old-35 or Old-64 gene or a portion thereof into the subject's cancerous cell under conditions permitting expression of the gene in the subject's cell so as to thereby reverse the cancerous phenotype of the cell.
- 25 30 27. The method according to claim 25 or 26, wherein the nucleic acid molecule comprises a vector.
- 35 28. The method according to claim 25 or 26, wherein the Old-35 or Old-64 gene is linked to a regulatory

-59-

element such that its expression is under the control of the regulatory element.

29. The method according to claim 26, wherein the
5 regulatory element is a tissue specific regulatory element.
30. The method of claim 25 or 26, wherein the nucleic acid
10 molecule is introduced into the cells by naked DNA technology, adenovirus vector, adeno-associated virus vector, Epstein-Barr virus vector, Herpes virus vector, attenuated HIV vector, retroviral vectors, vaccinia virus vector, liposomes, antibody-coated liposomes, mechanical or electrical means.
- 15 31. A method for reversing the cancerous phenotype of a cancer cell which comprises introducing OLD-35 or OLD-64 protein into the cancerous cell so as to thereby reverse the cancerous phenotype of the cell.
- 20 32. A method for reversing the cancerous phenotype of a cancer cell in a subject which comprises introducing OLD-35 or OLD-64 protein into the subject's cancerous cell so as to thereby reverse the cancerous phenotype
25 of the cell.
33. The method according to claim 25, 26, 31 or 32, wherein the cancer cell is a breast, cervical, colon, pancreatic, thyroid, skin, brain, prostate,
30 nasopharyngeal, lung, glioblastoma multiforme, lymphoma, leukemia, connective tissue, nervous system cell or basal cell.
34. A pharmaceutical composition which comprises an amount
35 of a nucleic acid molecule comprising Old-35, Old-64 gene or portion thereof effective to reverse the cancerous phenotype of a cancer cell and a

-60-

pharmaceutically acceptable carrier.

35. The pharmaceutical composition of claim 34, wherein the nucleic acid molecule comprises a vector.

5

36. The pharmaceutical composition of claim 35, wherein the vector is an adenovirus vector, adeno-associated virus vector, Epstein-Barr virus vector, Herpes virus vector, attenuated HIV vector, retrovirus vector or vaccinia virus vector.

10

37. A pharmaceutical composition comprising an amount of OLD-35 or OLD-64 protein effective to reverse the cancerous phenotype of a cancer cell and a pharmaceutically acceptable carrier.

15

38. The pharmaceutical composition of claim 34 or 36, wherein the cancer cell is a breast, cervical, colon, pancreatic, thyroid, skin, brain, prostate, nasopharyngeal, lung, glioblastoma multiforme, lymphoma, leukemia, connective tissue, nervous system or basal cell.

20

39. A method of determining whether a cell is senescent comprising measurement of the expression of the Old-35 gene, wherein the expression of the Old-35 gene indicates that the cell is senescent.

25

40. The method of claim 39, wherein the expression of the Old-35 gene is measured by the expression of Old-35 specific RNA.

30

41. The method of claim 39, wherein the expression of the Old-35 gene is measured by the expression of the OLD-35 protein.

35

-61-

42. A method of determining whether a cell is terminally differentiated comprising measurement of the expression of the Old-35 gene, wherein the expression of the Old-35 gene indicates that the cell is terminally differentiated.
43. The method of claim 42, wherein the expression of the Old- 35 gene is measured by the expression of Old-35 specific RNA.
44. The method of claim 42, wherein the expression of the Old-35 gene is measured by the expression of OLD-35 protein.
45. A method of determining whether a cell is growth arrested comprising measurement of the expression of the Old-35 gene, wherein the expression of the Old-35 gene indicates that the cell is growth arrested.
46. The method of claim 45, wherein the expression of the Old-35 gene is measured by the expression of old 35 specific RNA.
47. The method of claim 45, wherein the expression of the Old-35 gene is measured by the expression of OLD-35 protein.
48. A method of inhibiting growth of cancer cells comprising contacting the cancer cells with an amount of purified OLD-64 protein effective to inhibit growth of cancer cells.
49. A method of inhibiting growth of cancer cells comprising contacting the cancer cells with an amount of purified OLD-64 protein effective to inhibit growth of cancer cells.

-62-

50. A method of determining whether a cell is senescent comprising measurement of the expression of the Old-64 gene, wherein the expression of the Old-64 gene indicates that the cell is senescent.
- 5
51. The method of claim 50, wherein the expression of the old-64 gene is measured by the expression of Old-64 specific RNA.
- 10 52. The method of claim 50, wherein the expression of the Old-64 gene is measured by the expression of the OLD-64 protein.
- 15 53. A method of determining whether a cell is terminally differentiated comprising measurement of the expression of the Old-64 gene, wherein the expression of the Old-64 gene indicates that the cell is terminally differentiated.
- 20 54. The method of claim 53, wherein the expression of the Old-64 gene is measured by the expression of Old-64 specific RNA.
- 25 55. The method of claim 53, wherein the expression of the Old-64 gene is measured by the expression of the OLD-64 protein.
- 30 56. A method of determining whether a cell is growth arrested comprising measurement of the expression of the Old-64 gene, wherein the expression of the Old-64 gene indicates that the cell is growth arrested.
- 35 57. The method of claim 56, wherein the expression of the Old-64 gene is measured by the expression of Old-64 specific RNA.

-63-

58. The method of claim 56, wherein the expression of the Old-64 gene is measured by the expression of the OLD-64 protein.
- 5 59. A method of regenerating tissues comprising contacting the tissue with an inhibitor of OLD-35 or OLD-64 protein or a portion thereof at a concentration effective to regenerate said tissues.
- 10 60. A method of anti-aging in a cell comprising contacting the cell with an agent for inhibiting expression of Old-35 or Old-64 gene at a concentration effective to reverse the aging process in the cell.
- 15 61. A pharmaceutical composition for stimulating cell growth comprising a pharmaceutically acceptable carrier and purified Old-35 or Old-64 suppressant at a concentration effective to stimulate cell growth.
- 20 62. A method for screening the presence of interferon alpha or beta of a sample comprising steps of:
- (a) contacting the sample with cells under conditions permitting expression of Old-35 or Old-64 gene in the presence of interferon
25 alpha or beta; and
 - (b) determining the expression of the Old-35 or the Old-64 gene, an increase of expression indicates the presence of interferon alpha or beta.
- 30
63. A method for detection of the secretion of interferon alpha or beta comprising steps of:
- (a) obtaining an appropriate sample from the subject; and
 - 35 (b) detecting expression of Old-35 or Old-64 gene, the expression of the Old-35 or the

-64-

Old-64 gene indicating the secretion of interferon in a subject.

64. A method for monitoring chemotherapy of a subject
5 comprising steps of:
- (a) obtaining an appropriate sample from the subject; and
 - (b) detecting expression of Old-35 or Old-64
10 gene, the expression of Old-35 or Old-64 gene indicating that the chemotherapy is effective.
65. A method for diagnosis of the proliferating stage of a tumor from a subject comprising steps of:
- 15 (a) obtaining an appropriate sample from the subject; and
 - (b) detecting expression of the Old-35 or the Old-64 gene, the expression of the Old-35 or the Old-64 gene indicating that the tumor is
20 not at a proliferating stage.
66. A kit for diagnosis of the proliferating stage of a tumor, comprising a nucleic acid molecule capable of specifically hybridizing to the nucleic acid molecule
25 of the Old-35 or the Old-64 gene.
67. A kit for diagnosis of the proliferating stage of a tumor, comprising antibody capable of specifically recognizing OLD-35 or OLD-64 protein.
30
68. A method for identifying an agent that modulates the expression of the Old-35 or the Old-64 gene, comprising:
- 35 (a) contacting a candidate agent with a cell transformed or transfected with a reporter gene under the control of a Old-35 or Old-64

-65-

5 promoter or a regulatory element thereof
under conditions and for a time sufficient
to allow the candidate agent to directly or
indirectly alter expression of the promoter
or regulatory element thereof; and

10 (b) determining the effect of the candidate
agent on the level of reporter protein
produced by the cell, thereby identifying an
agent that modulates expression of Old-35 or
64 gene.

69. A method of identifying compounds that induce
proliferation or cancerous phenotype, comprising:
15 exposing cell comprising the promoter of Old-35 or
Old-64 to the compound and identifying compounds that
suppress the Old-35 or 64 promoter.

70. A method of identifying compounds that induces
senescence, or terminal differentiation, comprising:
20 exposing the cell comprising the promoter of Old-35 or
Old-64 to the compound and identifying compounds that
activate the Old-35 or 64 promoter.

71. A method of identifying genes which are common to the
25 pathway of senescence and terminal differentiation
comprising steps of:

- (a) obtaining a substrated library which is
enriched for genes expressed in terminal
differentiation;
- 30 (b) screening the library with senescent probe
to identify novel genes which are expressed
during senescence and terminal
differentiation; and
- 35 (c) examining the biological activity of the
identified gene to determined whether it is
expressed during senescence and terminal
differentiation.

-66-

72. A method of identifying genes which are common to the pathway of senescence and terminal differentiation comprising steps of:
- 5 (a) obtaining a subtracted library which is enriched for genes expressed in senescence;
 - (b) screening the library with terminal differentiation probe to identify novel genes which are expressed during senescence and terminal differentiation; and
 - 10 (c) examining the biological activity of the identified gene to determine whether it is expressed during senescence and terminal differentiation.
- 15 73. The gene identified by the method of claim 71 or 72.
74. A method of degrading specific RNAs in a cell comprising induction of the expression of Old-35 gene.
- 20 74. A method of degrading specific RNAs in a cell comprising introducing a vector into the cell comprising the Old-35 gene.
75. Expression of Old-35 can be used as diagnostic
25 indicator of cellular senescence, terminal differentiation and/or growth suppression.
- (a) can be used to determine if a cell has lost proliferative ability and become senescent.
- 30 76. Expression of Old-35 can be used as a marker to identify drugs or small molecules that will induce senescence, e.g., to inhibit cancer cell growth or abnormal proliferative states (such as psoriasis, hemangioblastoma, etc.)
- 35
77. Expression of Old-35 can be used to identify drugs or

-67-

small molecules that will inhibit senescence, possible uses including stimulating tissue regrowth, repair and regeneration.


- 5 78. Expression of Old-35 can be used as a marker to identify drugs or small molecules that will induce terminal cell differentiation, e.g., to inhibit cancer cell growth or abnormal proliferative states (such as psoriasis, hemangioblastoma, etc.).
- 10
79. Expression of Old-35 can be used to identify drugs or small molecules that will inhibit terminal differentiation, possible uses including stimulating tissue regrowth, repair and regeneration.
- 15
80. Expression of Old-35 can be used as marker for detecting cytokines, specifically type I interferons, in biological samples. Since type I interferon, including leukocyte and fibroblast interferons, which
- 20 activate gene expression through the well characterized Jak and Stat kinase pathways, this gene can be used to monitor for drugs and small molecules that activate these important pathways.
- 25 81. The combination of Old-35 with other interacting proteins can be used to target the differentiation of specific target cells. This can result in the reprogramming of pluripotent stem cells to terminally differentiated end cells.
- 30
82. Old-35 can be used to selectively stabilize specific mRNAs possibly containing AU rich 3' UTRs (untranslated regions). This effect can result in the sustained expression of genes potentiating or
- 35 inhibiting cell growth. It could also result in the stabilizing of cytokine genes resulting in increased biological and immunological activity.

-68-

83. Old-35 can be used as part of a methodology to polymerize random NTPs into nucleic acids.
- 5 84. Old-35 can be used to induce the degradation of specific mRNAs.

1/19

FIG. 1



2/19

FIG. 2A




FIG. 2B





FIG. 2C





FIG. 2D

EtBr


3/19

FIG. 3A

Human Multiple Tissue Northern Blot

**FIG. 3B**

Mouse Development



4/19

FIG. 4A


human	TTGAAGATTAC	AAATGGTGACATGGAC	CTTCAAAATAGCTGG	40
mouse	AAATGGTGACATGGAT	TTCAAAATAGCCGG	29
Consensus	aatggtgacatgga ttcaaaatagc gg			
human	CAC	TAATAAAGGAATAACT	TGCATTACAGGCTGATATTAAA	80
mouse	TAC	AAATAAAGGAATAACT	TGCATTACAGGCTGATATTAAAG	69
Consensus	ac aataaaggaataactgcattacaggctgatattaa			
human	TTACCTGGAA	TACCAAT	AAAAATTGTGATGGACGGCTATTC	120
mouse	TTACCTGGAG	TACCAAT	TAAAATTATAATGGAAAGCCATCC	109
Consensus	ttacctgga taccaat aaaatt t atgga gc at c			
human	AACAAGC	TCAGTGGCAAAA	AAGGAGATA	160
mouse	AACAAGC	TCAGTGGCAAAA	GAAGGAGATACTGCAGATAAT	149
Consensus	aacaagc tcagtggcaaaa aaggagata t cagat at			
human	GAACAAAAC	TATTTCAAAAACCTCGAGCATC	TAGAAAAGAA	200
mouse	GAACAAAAC	GATTTCAAAAACCTCGAGCATCA	AGAAAAGAA	189
Consensus	gaacaaaac atttcaaaaacctcgagcatc agaaaagaa			
human	AATGGACC	TGTTGTAGAAAC	TGTTCAAGGTTCCATTATCAA	240
mouse	AATGGACC	AGTTGTAGAAAC	AGTAAAGGTTCCATTATCAA	229
Consensus	aatggacc gttgtagaaac gt aggttccattatcaa			
human	AACGAGCAAAATT	TGTTGGACCTGGTGGC	TATAACTTAAA	280
mouse	AACGAGCAAAATT	CGTTGGACCTGGTGGATAT	CACTTAAA	269
Consensus	aacgagcaaaatt gttgg cctggtgg tat acttaaa			
human	AAAAC	TCAGGCTGAACAGGTTGTAAC	TATTAGTCAGGTG	320
mouse	AAAAC	TCAGGCTGAGACAGGTTGTAACA	AATTAGTCAGGTT	309
Consensus	aaaact caggctga acaggtgtaac attagtcaggt			
human	GATGAAGAAAC	GTTTCTGTG	TATTTGCACCAACACC	360
mouse	GATGAAGAAAC	CTTCTCCAT	TATTTGCACCAACACCT	349
Consensus	gatgaagaaac tt tc tatttgcaccaacacc a tg			
human	TTATGCATGA	GCAAGAGAC	TTTATTAC	400
mouse	CAATGCATGA	GCAAGAGAT	TTTATTAC	389
Consensus	atgcatga gcaagaga ttcattac gaaat tgca			
human	GGATGATCA	GAGCAGCAATT	AGAATTTGGAGCAGT	440
mouse	AGATGATCA	GAGCAACAATT	AGAATTTGGAGCAGT	429
Consensus	gatgatca gagca caattagaatttggagcagt tat			
human	ACCGC	ACAATAACTGAAATCAGAGA	TAATGGTGT	480
mouse	ACCGC	GACAATAACTGAAATCAGAGA	CACTGGAGTGATGG	469
Consensus	accgc acaataactgaaatcagaga actgg gt atgg			

5/19

FIG. 4B


human	TAAATTTATATCCAAATATGACTGCCGTACTGCTTCATAA	520
mouse	TAAACCTGTATCCAAACATGACTGCCAGTGCTGCTTCATAA	509
Consensus	taaaa t tatccaaa atgactgc gt ctgcttcataa	
human	CAACACAACCTTGAT.AACGAAAGATTAAACATCCTACTGCC	559
mouse	TTACACAACCTTGACC AACGAAAGATTAAACATCCCACTGCC	549
Consensus	cacaacttga aacgaaagattaaacatcc actgcc	
human	CTAGGATTAGAAAGTTGGCCAAGAAATTCAGGTCAAATACT	599
mouse	CTAGGACCTAGAGGTTGGCCAAGAAATTCAGGTCAAATACT	589
Consensus	ctagga taga gttggccaagaaattcaggt aaatact	
human	TTGGACCGTGACCCAGCCGATGGAAGAATGAGGCTTTCTCG	639
mouse	TTGGCCGTGATCCAGCTGATGGAAGAATGAGGCTTTCTCG	629
Consensus	ttgg cgtga ccagc gatggaagaatgaggctttctcg	
human	AAAAGTGCTTC	650
mouse	TAAAGTACTTC	640
Consensus	aaagt cttc	

6/19

FIG. 5

7/19

FIG. 6



8/19


FIG. 7

Hu GM-CSF	UAAU <u>AUUUA</u> UAU <u>AUUUA</u> UAUUUUAAAAU <u>AUUUA</u> UUU <u>AUUUA</u> UUUAA
Hu IFN- α	U <u>AUUUA</u> UUUA
Hu IL 2	U <u>AUUUA</u> UUUAAAUA <u>AUUUA</u> AAUUUUUAU <u>AUUUA</u> AU
Hu TNF	AAUUAA <u>AUUUA</u> UU <u>AUUUA</u> UUUAUU <u>AUUUA</u> UUUAUU
C-fos	GUUUUUAA <u>AUUUA</u> UUUAUUAAAGAUGGAUUCUCAGAU <u>AUUUA</u> UAUUUUUU AUUUUAUUUUUUUU
Old-35	A <u>UUUA</u> CAUGUGCCAUUUUUUUAAUUCGAGUAACCCAUAUUUGUUUAAUU GUA <u>AUUUA</u> CAUUAUAAAUCAAGAAAU <u>AUUUA</u> UUUAUUAAAAGUAAGUC <u>AUUUA</u> UACAUCUUAGA


9/19

FIG. 8A

Response of Old-35
To IFN- β Treatment
In the Presence of Cyclohexamide

**FIG. 8B**

Half-life of Old-35 in IFN- β +MEZ
Treated HO-1



10/19

FIG. 9A

GATGGTCCTT	TCCTTCTGCC	ACGGCGGGAT	CGGGCACTCA	CCCAGTTGCA
AGTGCGAGCA	CTATGGAGTA	GCGCAGGGTC	TCGAGCTGTG	GCCGTGGACT
TAGGCAACAG	GAAATTAGAA	ATATCTTCTG	GAAAGCTGGC	CAGATTTGCA
GATGGCTCTG	CTGTAGTACA	GTCAGGTGAC	ACTGCAGTAA	TGGTCACAGC
GGTCAGTAAA	ACAAAACCTT	CCCCTTCCCA	GTTTATGCCT	TTGGTGGTTG
ACTACAGACA	AAAAGCTGCT	GCAGCAGGTA	GAATTCCCAC	AAACTATCTG
AGAAGAGAGG	TTGGTACTTC	TGATAAAGAA	ATTCTAACAA	GTCGAATAAT
AGATCGTTCA	ATTAGACCGC	TCTTTCCAGC	TGGCTACTTC	TATGATACAC
AGGTTCTGTG	TAATCTGTTA	GCAGTAGATG	GTGTAAATGA	GCCTGATGTC
CTAGCAATTA	ATGGCGCTTC	CGTAGCCCTC	TCATTATCAG	ATATTCCTTG
GAATGGACCT	GTTGGGGCAG	TACGAATAGG	AATAATTGAT	GGAGAATATG
TTGTTAACCC	AACAAGAAAA	GAAATGTCTT	CTAGTACTTT	AAATTTAGTG
GTTGCTGGAG	CACCTAAAAG	TCAGATTGTC	ATGTTGGAAG	CCTCTGCAGA
GAACATTTTA	CAGCAGGACT	TTTGCCATGC	TATCAAAGTG	GGAGTGAAAT
ATACCCAACA	AATAATTCAG	GGCATTCCAGC	AGTTGGTAAA	AGAACTGGT
GTTACCAAGA	GGACACCTCA	GAAGTTATTT	ACCCCTTCGC	CAGAGATTGT
GAAATATACT	CATAAACTTG	CTATGGAGAG	ACTCTATGCA	GTTTTTACAG
ATTACGAGCA	TGACAAAGTT	TCCAGAGATG	AAGCTGTAA	CAAAATAAGA
TTAGATACGG	AGGAACAAC	AAAAGAAAA	TTCCAGAAG	CCGATCCATA
TGAAATAATA	GAATCCTTCA	ATGTTGTTGC	AAAGGAAGTT	TTTAGAAGTA
TTGTTTTGAA	TGAATACAAA	AGGTGCGATG	GTCGGGATTT	GACTTCACTT
AGGAATGTAA	GTTGTGAGGT	AGATATGTTT	AAAACCCCTC	ATGGATCAGC
ATTATTTCAA	AGAGGACAAA	CACAGGTGCT	TTGTACCGTT	ACATTTGATT
CATTAGAATC	TGGTATTAAG	TCAGATCAAG	TTATAACAGC	TATAAATGGG
ATAAAAGATA	AAAATTTTCA	GCTGCACTAC	GAGTTTCCTC	CTTATGCAAC
TAATGAAATT	GGCAAAGTCA	CTGGTTTAAA	TAGAAGAGAA	CTTGGGCATG
GTGCTCTTGC	TGAGAAAGCT	TTGTATCCTG	TTATTCCCAG	AGATTTTCCT
TTCACCATAA	GAGTTACATC	TGAAGTCTTA	GAGTCAAATG	GGTCATCTTC
TATGGCATCT	GCATGTGGCG	GAAGTTTAGC	ATTAATGGAT	TCAGGGGTTC
CAATTTTCATC	TGCTGTTGCA	GGCGTAGCAA	TAGGATTGGT	CACCAAAACC
GATCCTGAGA	AGGGTGAAAT	AGAAGATTAT	CGTTTGCTGA	CAGATATTTT
GGGAATTGAA	GATTACAATG	GTGACATGGA	CTTCAAAATA	GCTGGCACTA
ATAAAGGAAT	AACTGCATTA	CAGGCTGATA	TTAAATTACC	TGGAATACCA
ATAAAAATTG	TGATGGAGGC	TATTCAACAA	GCTTCAGTGG	CAAAAAAGGA
GATATTACAG	ATCATGAACA	AAACTATTTT	AAAACCTCGA	GCATCTAGAA
AAGAAAATGG	ACCTGTTGTA	GAAACTGTTT	AGGTTCCATT	ATCAAAACGA
GCAAAATTTG	TTGGACCTGG	TGGCTATAAC	TTAAAAAAC	TTCAGGCTGA
AACAGGTGTA	ACTATTAGTC	AGGTGGATGA	AGAAACGTTT	TCTGTATTTG
CACCAACACC	CAGTGTATATG	CATGAGGCAA	GAGACTTCAT	TACTGAAATC
TGCAAGGATG	ATCAGGAGCA	GCAATTAGAA	TTTGGAGCAG	TATATACCGC
CACAATAACT	GAAATCAGAG	ATACTGGTGT	AATGGTAAAA	TTATATCCAA
ATATGACTGC	GGTACTGCTT	CATAACACAC	AACTTGATAA	CGAAAGATTA
AACATCCTAC	TGCCCTAGGA	TTAGAAGTTG	GCCAAGAAAT	TCAGGTGAAA
TACTTTGGAC	GTGACCCAGC	CGATGGAAGA	ATGAGGCTTT	CTCGAAAAGT
GCTTCAGTCG	CCAGCTACAA	CCGTGGTCAG	AACTTTGAAT	GACAGAAGTA
GTATTGTAAT	GGGAGAACCT	ATTTACACAGT	CATCATCTAA	TTCTCAGTGA
TTTTTTTTTT	TTAAAGAGAA	TTCTAGAATT	CTATTTTGTC	TAGGGTGATG
TGCTGTAGAG	CAACATTTTA	GTAGATCTTC	CATTGTGTAG	ATTTCTATAT
AATATAAATA	CATTTTAATT	ATTTGTACTA	AAATGCTCAT	TTACATGTGC
CATTTTTTTA	ATTCGAGTAA	CCCATATTTG	TTTAATTGTA	TTTACATTAT
AAATCAAGAA	ATATTTATTA	<u>TTAAAAGTAA</u>	GTCATTTATA	CATCTTAGA

11/19

FIG. 9B

DGPFLPRRD RALTQLQVRA LWSSAGSRAV AVDLGNRKLE ISSGKLARFA
DGSVVQSGD TAVMVTAVSK TKPSPSQFMP LVVDYRQKAA AAGRIPTNYL
RREVGTSDE ILTSRIIDRS IRPLFPAGYF YDTQVLCNLL AVDGVNEPDV
LAINGASVAL SLSDIPWNGP VGAVRIGIID GEYVVPTRK EMSSSTLNLV
VAGAPKSQIV MLEASAENIL QQDFCHAIKV GVKYTOQIIQ GIQQLVKETG
VTKRTPQKLF TPSPEIVKYT HKLAMERLYA VFTDYEHDKV SRDEAVNKIR
LDTEEQLKEK FPEADPYEII ESFNVVAVEV FRSIVLNEYK RCDGRDLTSL
RNVSCVDMF KTLHGSALEF RGQTQVLCTV TFDSLESGIK SDQVITAING
IKDKNFMLHY EFPPYATNEI GKVTGLNRRE LGHGALAEKA LYPVIPRDFP
FTIRVTSEVL ESNGSSSMAS ACGGSLALMD SGVPISAVA GVAIGLVTKT
DPEKGEIEDY RLLTDILGIE DYNGDMDFKI AGTNKGITAL QADIKLPGIP
IKIVMEAIQQ ASVAKKEILQ IMNKTISKPR ASRKENGPPV ETVQVPLSKR
AKFVGPPGYN LKKLQAEAGV TISQVDEETF SVFAPTPSVM HEARDFITEI
CKDDQEQQLE FGAVYTATIT EIRDTGVMVK LYPNMTAVLL HNTQLDNERL
NILLP.

12/19

FIG. 10A

B subtilisMGQEKHVFTT	18
human	DGPFLPRRDRALTQLOVRALWSSAGSRAVA	40
Consensus	d r l	
B subtilis	VETGOLAKQANGAVMIRYGD TAVLS TATASKEPKPLDFF	58
human	ISSKILARFADGSAVVQSGDTAVMVTAVSKTKPS	80
Consensus	g l a a g gdtav ta p p f p	
B subtilis	ITVNEERLYAVGKIFGGFIKREGRPSENAVLASRLIDRP	98
human	LVVDYRQAAAAGRIP TNYLRREVGTSDEITTSRIIDRS	120
Consensus	l v y a g ip re s k l sr idr	
B subtilis	IRPLFADCFRNEVOI SIVMSVDONCSSEMAEMFGSSLAL	138
human	IRPLFPAGYFYDTQVLCNLLAVDGVNEPDVLAINGASVAL	160
Consensus	irplf g qv vd a g s al	
B subtilis	SVSDIPFEGHIACVTVSRIDDOFTIINPTVDQLEKSDINLV	178
human	SLSDIPWNGPVGAVRISIIDGEYVVNPTRKEMSSSTLNLV	200
Consensus	s sdip gp v g id npt s nlv	
B subtilis	VAGT.RDAINNEVAGADEVPPEEIMLEAIMFGHEEIKRLIA	217
human	VAGAPNSQIVLEASAENILOQDFCHAIKVGKVKYTOQIIQ	240
Consensus	vag k i m ea a ai g i	
B subtilis	FOEEIVAAVCKE.SEIKLFEIDEEELNEKVKALEEDLLK	256
human	GIOQLVKETGVTRTPOKLFTPSPEIVKYTHKLAMERLYA	280
Consensus	v g k klf e la e l	
B subtilis	AIQVEKHAREDAINEVKNNAVAKFEDEHDEDTIKQVKQ	296
human	VFTDYEHDKVSRDEAVNKIRLDTEEQLKPKFPEADPYEII	320
Consensus	e k e	
B subtilis	ILSKLVKNEVRRLITE.EKVRPDGRGVDQIRPLSSEVGLL	335
human	ESFNVVAKEVFRSIVLNLYKRC DGRDLTSIRNVSCVDMF	360
Consensus	v ev r i e r dgr r s ev	
B subtilis	PRTHGSGLETRGOTCALSVCTLGALGDVQILDGLGVEES.	374
human	KTLHGSALFQRCOTOVLCTVTFDSIESGIKSDQVITAING	400
Consensus	hgs lf rgqtq l t l d	
B subtilis	...KRFMHYNYFPQFSVGETGPMRGPCRREIGHGALGERA	411
human	IKDKNEHLYEFPYPATNEIGKVTGLNRRELGHGALA EKA	440
Consensus	k fm hy fp e g g rre ghgal e a	
B subtilis	LEPVIPSEKDFPYTVRLVSEVLESNGSTSASICASTLAM	451
human	LYPVIPR...DFPFTIRVTSEVLESNGSSMASACGGSLAL	478
Consensus	l pvip dfp t r sevlesngs s as c la	

13/19

FIG. 10B

B subtilis	MDAGVPIKAPVAGIAMGLVKS.....EHYTVLTDIQG	484
human	MDSCVPISSAVAGVAIGLVTKTDPEKGEIEDYRLTDLG	518
Consensus	md gvpi vag a glv e y ltdi g	
B subtilis	MEDALGDMDFRVAGTEKGVLTALQMDIKIEGLSREILLEAL	524
human	IEDYNGDMDFRIAGTNKGITALQADIKLPCIPIKIVMEAI	558
Consensus	ed gdmfdk agt kg talq dik g i ea	
B subtilis	QQAQKGRMEIINSLATLSESRKELSRYPKILMTINPD	564
human	QQAQSVAKREILOINNKTIISKPRASRKENGIVVETVQVPLS	598
Consensus	qqa eil m t s r p t	
B subtilis	KIRDVISESKQINHLIEETGVKIDIEQDGTIFISSTDES	604
human	KRAKFVCGGYNLKKLOAETGVLSQVDEETFSVFAPTPS	638
Consensus	k gp g k etgv i t s	
B subtilis	GNQKPKKILIEDLVREVEVGQLYLSKVVKRIEKFGAFVEIFS	644
human	VMHEARDFTTEICKDDQEQQLFEFGAVYTATITEIRD TGVM	678
Consensus	a i ql g v	
B subtilis	GNQGLVHISEIALERVGVKEDVVKIGDEILVKVTEIDKQG	684
human	VNLYPNMTAVLLENTQLDNERLNILLP.....	705
Consensus	k l e	
B subtilis	RVNLSRKAVLREEKEKEEQQS	705
human	705
Consensus		

Half-life of Old-35 mRNA




FIGURE 11

FIGURE 12




FIGURE 13




FIGURE 14

The effect of subtypes of IFN- α on
Old-35 expression




FIGURE 15

Old-35 is expressed in the spinal column
and the genital area



FIGURE 16

Localization of Old-35 In HeLa cells



GFP
100X

GFP-hPNPase
40X

SEQUENCE LISTING

<110> Fisher, Paul B.

<120> Genes Displaying Enhanced Expression During
Cellular Senescence and Terminal Cell
Differentiation and Uses Thereof

<130> 0575/56765

<140> WIPO ST. 10/C

<141> 1999-02-03

<160> 50

<170> PatentIn Ver. 2.0

<210> 1

<211> 674

<212> DNA

<213> Homo sapien

<400> 1

```

aattcggcac gagcacgtct tgacctgaa cgcaaagtgg aatctttgca agaagagatt 60
gcctttttga agaaactcca cgaagaggaa atccaggagc tgcaggctca gattcaggaa 120
cagcatgtcc aaatcgatgt ggatgtttcc aagcctgacc tcacggctgc cctgcgtgac 180
gtacgtcagc aatatgaaag tgtggctgcc aagaacctgc aggaggcaga agaatggtac 240
aaatccaagt ttgctgacct ctctgaggct gccaacccga acaatgacgc cctgcgccag 300
gcaaagcagg agtccactga gtaccggaga cagggtgcagt ccctcacctg tgaagtggat 360
gcccttaaag gaaccaatga gtccctggaa cgccagatgc gtgaaatgga agagaacttt 420
gccgttgaag ctgctaacta ccaagacact attggcccgc ctgcaggatg agattcagaa 480
tatgaaggag gaaatggctc gtcaccttcg tgaataccaa gacctgctca atgntaagat 540
ggcccttgac attgagattg ccacctacag gaagctgctg ggaaggcgan gagagcagga 600
tttctctgct cttccaaact tttcctcctt gaccttgagg gaaactaatc tggattcact 660
ccctcttggg tgaa 674

```

<210> 2

<211> 678

<212> DNA

<213> Homo sapien

<400> 2

```

aattcggcac gagcaggacc caaggaacca aaattgcac tgatgggtctc aagggtcgtg 60
tgtttgaagt gagtcttgct gatttgcaga atgatgaagt tgcattagaa aattcaagct 120
gattactgaa gatgttcagg gtaaaaaactg cctgactaac ttccatggca tggatcttac 180
ccgtgacaaa atgtgttcca tggtcaaaaa atggcagaca atgattgaag ctcacgttga 240
tgtcaagact accgatgggt acttgcttcg tctgttctgt gttgggttta ctaaaaaacg 300
caacaatcag atacggaaga cctcttatgc tcagcaccaa cagggtccgc aaatccggaa 360

```

```

gaagatgatg gaaatcatga cccgagaggt gcagacaaat gacttgaaag aagtgggtcaa 420
taaattgatt ccagacagca ttggaaaaga catagaaaag gcttgccaat ctatttatcc 480
tctccatgat gtcttcgtta gaaaagtaaa aatgctgaag aagcccaagt ttgaattggg 540
aaagctcatg gagcttcatg gtgaanggca gtagttctgg aaaaagccac ttggggacga 600
aacagggtgct aaaagtttga acgactgatg gatattgaac cccagtccaa gaatctgggt 660
aaaggtcana cttcaaat
678

```

<210> 3

<211> 670

<212> DNA

<213> Homo sapien

<400> 3

```

aattcggcac gaggatgatg accttcaaga aaatgaagac aataaacaac ataaagaaag 60
cttgaaaaga gtgaccttgg ctttaccaga tgatgcggaa actgaagata cagggtgtttt 120
aaatgtaaag aaaaattctg atgaagttaa atcctccttt gaaaaaagac aggaaaagat 180
gaatgaaaaa attgcatctt tagaaaaaga gttgttagaa aaaaagccgt ggcaacttca 240
gggggaagtg acagcacaga agaggccaga gaacagcctc ctggaggaga ccctacactt 300
tgaccatgct gtccggatgg cacctgtgat tacagaggaa accacccttc aactggaaga 360
tatcattaaa cagaggataa gagatcaggc ttgggatgat gtagtacgta aagaaaaacc 420
taaagaggat gcatatgaat ataaaaagcg tttaacctta gaccatgaga agagtaaatt 480
gagccttgct gaaatttatg aacaggagta catcaaactc aaccagcaaa aaacagcaga 540
agaagaaaat ccagaacatg tagaaattca gaagatgatg gattccctct tcttaaattg 600
gatgcctctc aaacttccct ttatccctta accgcctgtc cagagattaa agttgnggcc 660
aaatctgcca
670

```

<210> 4

<211> 675

<212> DNA

<213> Homo sapien

<400> 4

```

aattcggcac gagatctgct gcaagcagcc aaatgttatg agaaggaact gggccgcctg 60
ctaagggatg ccccttcagg cataggcagt attttcctgt cagcatctga gcttgaggat 120
ggtagtgagg aaatgggcca gggcgagtc agctccagtc ccagagagct cctctctaac 180
tcagagcaac tgaactgaga cagaggagga aaacagagca tcagaagcct gcagtgggtg 240
ttgtgacggg taggaggata ggaagacagg gggccccaac ctgggattgc tgagcaggga 300
agctttgcat gttgctctaa ggtacatttt taaagagttg ttttttggcc gggcgagtg 360
gctcatgcct gtaatcccag cactttggga ggccgaggtg ggccggtcac gaggtctgga 420
gtttgagacc atcctggcta acacagtga atcccgcttc tactaaaaat acaaaaaatt 480
agccaggcgt ggtggtggc acctgtagtc ccagctactt gggagctgan gcangagaat 540
ggcgtgaacc tggaaggaa aagttgcagg tgagcccaag attgcgcccc cttgcactcc 600
agctgggcaa cagagcaaga cttcatctca aaaaaaaaaa aaaaaaactn ncggnggggg 660
gccccgggc cccca
675

```

<210> 5

<211> 460

<212> DNA

<213> Homo sapien

<400> 5

```
aattcggcac gagcacctct gtgtctacca tgacccccctt cctcacactg acctgtgttc 60
cttcctgtgt ctcttttcta ttaaaaataa gaacctgggc agagtgcggc agctcatgcc 120
tgtaatccca gcacttaggg aggccgagga gggcagatca cgaggtcagg agatcgaac 180
catcctggct aacacgggtga aaccccgctct ctactaaaaa atacaaaaaa ttagctgggc 240
gcagaggcac gggcctgtag tcccagctac tcaggaggcg gaggcaggag aatggcgta 300
acccgggagg cggagggtgc agtgagccag gattgtgcga ctgcactcca gcctgggtga 360
cagggtgaaa cgccatctca aaaaataaaa attaaaaaaa aaaaaaaaaa nntcnngggg 420
ggncccggtg ccnatttcnc cntatnggga gncntnncaa 460
```

<210> 6

<211> 445

<212> DNA

<213> Homo sapien

<400> 6

```
aattcggcac gagttctgcc catgctgcag acagtggcca agaacaagga ccagggcacc 60
tatgaggatt atgtcgaagg acttcgggtg tttgacaagg aaggaaatgg caccgtcatg 120
ggtgctgaaa tccggcatgt tcttgtcaca ctgggtgaga agatgacaga ggaagaagta 180
gagatgctgg tggcagggca tgaggacagc aatggttgta tcaactatga agagctcgtc 240
cgcattggtg tgaatggctg aggaccttcc cagtctcccc agagtccgtg cttttccctg 300
tgtgaatttt gtatctagcc taaagtttcc ctaggctttc ttgtctcagc aactttccca 360
tcttgtctct cttggatgat gtttgccgtc agcattcacc aaataaactt gctctctggg 420
ccctcggaaa aaaaaaaaaa aaaaa 445
```

<210> 7

<211> 666

<212> DNA

<213> Homo sapien

<400> 7

```
aattcggcac gaggcaatgt gcttggtttt aaagaaattc tccttgggaa aaaagtatcc 60
tcttttaatt ttacttccca taagcgtaaa tgacctggaca tagctcttgt gcaaccttta 120
aataaattgt tttgagtgtt ttttgagccc cagacaaata atgtttttaa gttatcccc 180
tgctacttta ctgatacctt tatcattcct gagacagttt gctaatttaa aaatgtagca 240
ttccatttgt atttatttct ctcccttgcc aaaaagattt tctaatactg cttgtaccag 300
ccagagaaag atccaaaaca ctactcagct ctcttgcaact gaggaattt ttccccctac 360
attgaactcct ggccatcac agccaaactt aaccttggtg gggtttggtt ttgatagcca 420
attagtcttg tgctggttgc aaagaattga tatttagatg gtttttaata ctacagagat 480
tgncttctct tatattngt cttttttatg ttgcatgttg cttttgntat cagcctgatt 540
ttttgctcag tatatgatag ttctgtgat ggtttgggta ttgggcagac atatcttcat 600
taagagtttt tggaaaactc atcaaattcg atgaatacat tttcttcata acccattgga 660
aatatc 666
```

<210> 8

<211> 409

<212> DNA

<213> Homo sapien

<400> 8

```

aattcggcac gagcgactac ggcgactaa tcttcaactc ctacatactt cccccattat 60
tcctagaacc aggcgacctg cgactccttg acgttgacaa tcgagtagta ctcccgattg 120
aagcccccat tcgtataata attacatcac aagacgtctt gcactcatga gctgtcccca 180
cattaggctt aaaaacagat gcaattcccg gacgtctaaa ccaaaccact ttcaccgcta 240
cacgaccggg ggtatactac ggtcaatgct ctgaaatctg tggagcaaac cacagtttca 300
tgcccatcgt cctagaatta attcccctaa aaatctttga aatagggccc gtattttacc 360
tatagcacc cctctacccc ctctagagca aaaaaaaaaa aaaaaaaaaa 409

```

<210> 9

<211> 667

<212> DNA

<213> Homo sapien

<400> 9

```

aattcggcac gagacaaggc acaaatgatc caaatcaagg aggctacaaa agggcagcct 60
agagggcgaga acagagaaaa gctagacaaa atgataagat cagccatatt tcattttgaa 120
tctgcagtgg aaaaaaagcc cacatttgag gtggctcatc tagacctggc aagaatgtat 180
atagaagcag gcaatcacag aaaagctgaa gagaattttc aaaaattggt atgcatgaaa 240
ccagtggtag aagaaacaat gcaagacata catttccact atggctcggt tcaggaattt 300
caaaagaaat ctgacgtcaa tgcaattatc cattatttaa aagctataaa aatagaacag 360
gcatcattaa caagggataa aagtatcaat tctttgaaga aattggtttt aaggaaactt 420
cggagaaagg cattanactg gaaagcttga gcctccttgg gttcgtctac aaattggaag 480
gaaatatgaa tgaagccctg gagtactatg agcgggccct gagactggct gctgactttg 540
agactctgtg agacaaggtc cttagcccca gatatcagcc ctttccattt catttcattt 600
tatgctaaca tttactaatc atcttttctg cttactgggt tcagaacctt ataattccct 660
ggnatga 667

```

<210> 10

<211> 672

<212> DNA

<213> Homo sapien

<400> 10

```

aattcttcct gtacgattgg ggatataacg ggcttcacta accttcccta ggcattgaaa 60
cttcccccaa atctgatgga cctagaagtc tgcttttgta cctgctgggc cccaaagttg 120
ggcatttttc tctctgttcc ctctcttttg aaaatgtaaa ataaaaccaa aaatagacaa 180
ctttttcttc agccattcca gcatagagaa caaaccttat ggaaacagga atgtcaattg 240
tgtaatcatt gttctaatta ggtaaataga agtccttatg tatgtgttac aagaatttcc 300
cccacaacat cctttatgac tgaagtcaa tgacagtttg tgtttgggtg taaaggattt 360
tctccatggc ctgaattaag accattagaa agcaccaggc cgtgggagca gtgaccatct 420
gctgactggt cttgtggatc ttgtgtccag ggacatgggg tgacatgcct cgtatgtgtt 480
agaggggtgga atggatgtgt ttggcgctgc atgggatctg gtgccccctc tctcctggat 540
tcacatcccc acccagggcc cggttttact aagtgtctgc cctanattgg gtcaaaggag 600
gtcatccaac tgactttatc aagtggaatt gggatatatt tgatatactt ctggctaaca 660

```

acatgggaaa ag

672

<210> 11

<211> 672

<212> DNA

<213> Homo sapien

<400> 11

```

aattcttcct gtacgattgg ggatataacg ggcttcacta accttccta ggcattgaaa 60
cttcccccaa atctgatgga cctagaagtc tgcttttgta cctgctgggc cccaaagtgg 120
ggcatttttc tctctgttcc ctctcttttg aaaatgtaaa ataaaaccaa aaatagacaa 180
ctttttcttc agccattcca gcatagagaa caaaccttat ggaaacagga atgtcaattg 240
tgtaatcatt gttctaatta ggtaaataga agtccttatg tatgtgttac aagaatttcc 300
cccacaacat cctttatgac tgaagttaa tgacagtgtg tgtttggtgg taaaggattt 360
tctccatggc ctgaattaag accattagaa agcaccaggc cgtgggagca gtgaccatct 420
gctgactgtt cttgtggatc ttgtgtccag ggacatgggg tgacatgcct cgtatgtgtt 480
agagggtgga atggatgtgt ttggcgctgc atgggatctg gtgcccctct tctcctggat 540
tcacatcccc acccagggcc cggttttact aagtgtctgc cctanattgg gtcaaaggag 600
gtcatccaac tgactttatc aagtggaatt gggatatatt tgatatactt ctggctaaca 660
acatgggaaa ag

```

672

<210> 12

<211> 669

<212> DNA

<213> Homo sapien

<400> 12

```

aattcctaga caccaaatac agtgtgggaa tacacaacct actagcctat gtgaaacacc 60
tgaaaggcca gaatgaggaa gccctgaaga gcttaaaaga agctgaaaac ttaatgcagg 120
aagaacatga caaccaagca aatgtgagga gtctgggtgac ctggggcaac tttgcctgga 180
tgtattacca catgggcaga ctggcagaag ccagactta cctggacaag gtggagaaca 240
tttgcaagaa gctttcaaat cccttccgct atagaatgga gtgtccagaa atagactgtg 300
aggaaggatg ggccttgctg aagtgtggag gaaagaatta tgaacggggc aaggcctgct 360
ttgaaaaggt gcttgaagtg gaccctgaaa accctgaatc cagcgctggg tatgcatct 420
ctgcctatcg cctggatggc tttaaattag ccacaaaaaa tcacaagcca ttttctttgc 480
ttccccaaag gcaggctgtc cgcttaaatc cagataatgg atatattaag ggtctccttg 540
ccctgaagct tcaggatgaa ggacaggaaa cttgaaggag aaaagtnat tgaanaactn 600
taccacccat gtccctcaga cctatgcttt gattgcagcc aagttttacc gaaaaaaagn 660
tntgggata

```

669

<210> 13

<211> 702

<212> DNA

<213> Homo sapien

<400> 13

```

aattcggcac gagtgttata ggagatcaca aatcaacatc tcacttccga accggggaag 60
aagacaagaa aattaatgaa gaactggagt ctcaatatca gcaaagtatg gacagtaa 120

```

```

tattcaggaag atatcggcga cattgtggac ttggcttcag tgaggtagaa gaccatgatg 180
gagaagggtga tgtggctgga gatgatgatg atgacgatga tgattcacct gatcctgaaa 240
gtccagatga ttctgaaagc gattcagagt cagagaaaga agaattctgct gaagaactcc 300
aagctgctga gcaccctgat gaagtggagg atcccaaaaa caaaaaagat gcaaaaagca 360
attataaaat gatgtttgtt aaatccagtg gttcataact cccaaacgct tagtctttgt 420
attaaaagta agccttattg ttacaatgca cagtggagga ctgcttatag agcacagacc 480
tttgtattat aattttttaa aaggcccttt taaataatta caaagagtgn ttgctttcaa 540
atgccatggg ttacactttt atgggcatga ctataccatt ttgnaaaga gtagagttgn 600
ataaaaataag aaatanntcc agtactcact tccttctatt agcatctcac cctntaatte 660
ccttatgggg aaatgcttct tttggttggg atagcttttt an 702

```

<210> 14

<211> 312

<212> DNA

<213> Homo sapien

<400> 14

```

aattcggcac gaggtaaatg ttgagccttt ttggcctggc ttgtttgcaa aggccttggc 60
caacgtcaac attgggagcc tcactgcaa tgtaggggcc ggtggacctg ctccagcagc 120
tggtgctgca ccagcaggag gtccctcccc ctccactgct gctgctccag ctgaggagaa 180
gaaagtggaa gcaaagaaag aagaatccga ggagtctgat gatgacatgg gcttttggct 240
ttttgactaa acctctttta taacatgttc aataaaaagc tgaactttaa aaaaaaaaaa 300
aaaaaaaaaa ac 312

```

<210> 15

<211> 391

<212> DNA

<213> Homo sapien

<400> 15

```

aattctgagg aggaactgat gatggcatgg aagaactttt cagtcatctg aaggaggttc 60
cgctggtttt cctcaaggct ctctgatggt tctaacctgg taggatccac ttcaaagcta 120
acatgttgcc aatcagagga tgtgatcaca attcgtaata aaggatccag gagtttttgt 180
agataggtag caccatatac cttgaaacag aatgtcatta ttttactggc caagctgttg 240
cctcggaaga gagtctgcat ggagtctgcc aattctactt ctttagaaaa catgttccag 300
agcagttggg agagtaaag ccgagaatca aacagagtaa ccagaactcg agggggggcc 360
cggtagccaa ttcgccctat agtgagtcgt t 391

```

<210> 16

<211> 720

<212> DNA

<213> Homo sapien

<400> 16

```

aattcnnatc gagnttcttt gtatgtgaac ggtcctggaa ggattctgtt gtccttggc 60
aggtgtgtgg ttgtcgctat agactggctc cggatgatctg gccattatac tctgctgtct 120
ccatcttgag gatgtagggg attatgctgt ctatcgaaac attgccaatg agaccagtaa 180
aaaaaagttc ttctgttatg ttggagctca tcagcctgag tgccggcagg cgaacagga 240

```


tccggggccaa tctataaaaag ggagtgtcat tagaaaagga gactgtttga tggccttcaa 300
 ccacagctca gcaaaggctc ctgggggtccc gtctgtattg caccagaatc aaaccaacng 360
 gatccacctt ccaccacact ttnttttctg atttcaacag ttctctttat agaaatttat 420
 catgagaaaa aaccaaataa gtccaaaang tatgtncana tgggttccct tcnctctggg 480
 aatccaactt tcttaacccc nccnccaaaa aaaanctngg aattcttnac cngngngnca 540
 ccttaaggng gaagccttca tnggaannac ttgctanana ctcattttaa aaaccgatta 600
 ntccaacccc tgtnttttct gnccnggaa aanacntccc ntgacatatg gctcaaataa 660
 aagggttttaa aggggaantt ttnaaaaaaa anaaaaaaa aaaccctngg gggggggccc 720

<210> 17

<211> 205

<212> DNA

<213> Homo sapien

<400> 17

aattcgaaca gcataacccc gattccgcta cgaccaactc atacacctcc tatgaaaaaa 60
 cttcctacca ctcaccctag cattacttat atgatatgtc tccataccca ttacaatctc 120
 cagcattccc cctcaaactt aaaaaaaaaa aaaaaaannt ngnggggggg cccggncccc 180
 anttcnccnt ntngggngnn gnntt 205

<210> 18

<211> 691

<212> DNA

<213> Homo sapien

<400> 18

aattcttaca tgttttcttt gctttaagt taactggcag ttttccattg gtttacctgt 60
 gaaatagttc aaagccaagt ttatatataa ttatatcagt cctctttcaa aggtagccat 120
 catggatctg gtagggggaa aatgtgtatt ttattacatc tttcacattg gctattttaa 180
 gacaaagaca aattctgttt cttgagaaga gaattattagc tttactgttt gttatggctt 240
 aatgacacta gctaataatca atagaaggat gtacatttcc aaattcacaa gttgtgtttg 300
 atatccaaag ctgaatacat tctgctttca tcttggtcac atacaattat ttttacagtt 360
 ctcccaaggg agttaggcta ttcacaacca ctcatcaca agtgaaatt aaccatagat 420
 gtagataaac tcagaaattt aattcatgtt tcttaaattg gctactttgt cctttttgnt 480
 attagggtgg tatttagtct attagccaca aaattgggaa aggagtagaa aaagcagtaa 540
 ctgacaactt gaataatata ccagagataa tatgagaatc agatcatttc aaaactcatt 600
 tcttatgtaa ctgcattgag aactgcatat gtttcgctga tatatggggg tttccatttg 660
 cgaatgggtc cattctctct ccggactttt t 691

<210> 19

<211> 483

<212> DNA

<213> Homo sapien

<400> 19

tctagaacta gtggatcccc cgggctgcag gaattcggca cgaggtttta agtactctga 60
 aattgatctg tgatcaataa tactaatatg ttatctttta ccgtattctg cctctcacta 120
 ttgattttta ttagtttaga gtatttgagc tggtatttct tgagcttaat attttttttag 180

```

agttaactct ttaaggagat aatcatggct gtagacaagg ccagggctgg ctgacgtgcc 240
ttagaaagtt tgaatgcaat aaagcgggtg ttggcggttct cctgcattgt agtgcggggt 300
acaaatgcta attgttccgt caactgggtg cagcagatga gccgcccact acagacgggt 360
actgcccagg gacctgcccc ggccccaccc aaggggtccc aaggggttgag atttctgcag 420
acctatagcc agcacactta gtcctgccct atatagagtt cctcttcggg aagcttttga 480
taa 483

```

<210> 20

<211> 589

<212> DNA

<213> Homo sapien

<400> 20

```

gcacgagtcg aaatgtacat tgggtgattct gaagcttata tcggagcaga cattaagac 60
aaattaaaaat gttatgactt tgatgtgcat acaatgaaga cactaaaaaa cattatttca 120
cctccgtggg atttcaggga atttgaagta gaaaaacaga ctgcagaaga aacgggggtt 180
acgccattgg aaacctcaag gaaaactcca gattccagac cttccttgga agaaaccttt 240
gaaattgaaa tgaatgaaag tgacatgatg ttagagacat ctatgtcaga ccacagcacg 300
tgactccagt cagtggctct gggtccactg tcccagtgtg ggtagtatt ccttcacatc 360
ctctccatgg cttaagaatg tcccacttcc taactgtgact ccaaactgca tctctacatt 420
taggaacaga gacccgcctt aagagactgg atcgcacacc ttgcaacag atgtgttctg 480
attctctgaa cctacaaaat agttatacat agtggaataa agaaggtaaa ccatcaaaaa 540
aaaaaaaaaa aaaccnncngg gggggcccg gccaatttg cccttangg 589

```

<210> 21

<211> 713

<212> DNA

<213> Homo sapien

<400> 21

```

aattcaagtg cctgattaat tgaggtggca acatagtttg agacgagggc agagaacagg 60
aagatacata gctagaagcg acgggtacaa aaagcaatgt gtacaagaag actttcagca 120
agtatacaga gagttcacct ctactctgcc ctccatcatag tcataatgta gcaagtaaag 180
aatgagaatg gattctgtac aatacactag aaaccaacat aatgtatttc tttaaaacct 240
gtgtgaaaaa ataaatgttc caccagtagg gataggggaa aagtaacca aagagagaaa 300
gagaaaggaa tgctgggttta tctttgtaga ttgtaatcga atggagaaat ttgcagtatt 360
ttagccacta ttaggaattt ttttttttng taaaangaan actgaactnt gttcaaaangc 420
tttcatganc ctgggttgaa acggtaggaa agcaccaaaa cgngggancc tggggactaa 480
gggcctgggt caaggacttg ggaaatggca ttgataatan atgggggggt tttccccct 540
ttaaaaatgt tggatnttaa gggatataac ccttntttta ctccgaaaat nttntgagaa 600
atcccaaaat tcncgggatg cttggaacca ttganatttt ntaggggaan gccttgaata 660
gcctanacct caaagttggn gngaacaaaa attggagccn ttgncccacc tcc 713

```

<210> 22

<211> 480

<212> DNA

<213> Homo sapien

<400> 22

cggcacgaga agaagtggta caggaggaat ttgtgatgat gagctgatct taatcaaaaa 60
 tactaaggct cgtacgtctg catcgattat cttacgtggg gcaaatgatt tcatgtgtga 120
 tgagatggag cgctctttac atgatgcact ttgtgtagtg aagagagttt tggagtcaaa 180
 atctgtgggt cccgggtggg gtgctgtaga agcagccctt tccatatacc ttgaaaacta 240
 tgcaaccagc atgggggtctc gggaacagct tgcgattgca gagtttgcaa gatcacttct 300
 tgttattccc aatacactag cagttaatgc tgcccaggac tccacagatc tggttgcaa 360
 attaagagct tttcataatg aggccaggt taaccagaa cgtaaaaatc taaaatgatt 420
 ggtcttgatt tgagcaatgg taaacctcga gggggggccc ggtacccaat tcgccctata 480

<210> 23

<211> 198

<212> DNA

<213> Homo sapien

<400> 23

cctgttaaaa gctgttcttg nggtttacat gtaacagaca tggtaaatat ttgtttacag 60
 tctttgttta acaaaccatg catttaagtt taagtgaagt caacaaaaag gaaatagggtg 120
 tatggatatg tgattttgag attaaagtta gtcttaaaat gtaaaaaaaaa aaaaaaaaaa 180
 aaaaaaaaaa aaaaaaaaaa 198

<210> 24

<211> 414

<212> DNA

<213> Homo sapien

<400> 24

aattcggcac gagaaaagca gtataactgc ctgacacagc gggattgaac gagagaagaa 60
 attgttcggt attgttcaga aaattcaaac acgcaaagat cttatggata aaactcagaa 120
 agtgaagggtg aagaaagaaa cggatgaactc cccagctatt tataaatttc agagtcgctc 180
 aaaacggtga cgtgttatag ataagccttg tcattctgta tcaaaaatct gttgtcgttt 240
 tctagtaact tcaaattcca ttactccaaa tggcatgggt ttccgggttg taaccataac 300
 taaattgtca gtctgacatt taatgtcttt ctatggacaa cattaaatct ccctcccttc 360
 tgtagaanan anannnnaaa aancncncg gggggggccg ggtccccatt cccc 414

<210> 25

<211> 367

<212> DNA

<213> Homo sapien

<400> 25

aattcggcac gagaaaagca gtataactgc ctgacacagc gggattgaac gagagaagaa 60
 attgttcggt attgttcaga aaattcaaac acgcaaagat cttatggata aaactcagaa 120
 agtgaagggtg aagaaagaaa cggatgaactc cccagctatt tataaatttc agagtcgctc 180
 aaaacggtga cgtgttatag ataagccttg tcattctgta tcaaaaatct gttgtcgttt 240
 tctagtaact tcaaattcca ttactccaaa tggcatgggt ttccgggttg taaccataac 300
 taaattgtca gtctgacatt taatgtcttt ctatgggaca acattaaatc tccctccctt 360
 ctgtaaa 367

<210> 26
 <211> 432
 <212> DNA
 <213> Homo sapien

<400> 26
 aattcggcac gaggcagact tgaacagtt ctgtctgcag aatgctcaac atgaccctct 60
 gctgactgga gtatcttcaa gtacaaatcc cttcagaccc cagaaagtct gtcccttttt 120
 gtagtaaaat gaatctttca aaggtttccc aaaccactcc ttatgatcca gtgaatattc 180
 aagagagcta catcttgaagc ctgtacaaaa gcttatccct gtaacacatg tgccataata 240
 tacaaacttc tactttcgtc agtccttaac atctacctct ctgaattttc atgaatttct 300
 atttcacaag ggtaattggt ttatatacac tggcagcagc atacaataaa acttagtatg 360
 aaactttaaa aaaaaaaaaa aaaacntcnn ggggggnccc ggancccant tcnccntata 420
 gggngnccgn tt 432

<210> 27
 <211> 398
 <212> DNA
 <213> Homo sapien

<400> 27
 aattcggcac gaggacaaaa ccagttggtg gtgacaagaa cggcgggtacc cgggtgggta 60
 aacttcgcaa aatgcctaga tattatccta ctgaagatgt gcctcgaaag ctgttgagcc 120
 acggcaaaaa acccttcagt cagcacgtga gaaaactgag agccagcatt acccccggga 180
 ccattctgat catcctcact ggacgccaca ggggcaagag ggtgggtttc ctgaagcagc 240
 tggctagtgg cttattactt gtgactggac ctctggctct caatcgantt cctctacnaa 300
 gaacacacca gaaatttgtc attgccactt caacccaaat cgatntcngc antgtannaa 360
 atcccaanac atcttactga tgcttacttc aagatgaa 398

<210> 28
 <211> 232
 <212> DNA
 <213> Homo sapien

<400> 28
 aattcggcac gagattgtat cggttttata ttacctgttc tgcttcacca ggagatcatg 60
 ctgctgtgat actgagtttt ctaaacagca taaggaagac ttgctcccct gtcctatgaa 120
 agagaatagt tttggagggg agaagtggga caaaaaagat gcagttttcc tttgtattgg 180
 gaaatgtgaa aataaaattg tcaactcttt caaaaaaaaaa aaaaaaaaaa aa 232

<210> 29
 <211> 539
 <212> DNA
 <213> Homo sapien

<400> 29
 aattcggcac gagcacaacc agaaagtaag gtgttctact tgaaaatgaa aggagattat 60

```

tttaggtatc tttctgaagt ggcattctgga gacaacaaac aaaccactgt gtcgaactcc 120
cagcaggctt accaggaagc atttgaaatt agtaagaaag aaatgcagcc tacacaccca 180
attcgtcttg gtctggcact aaatctctca gtcttttact atgagattct aaactctcct 240
gaaaaggcct gtagcctggc aaaaacggca tttgatgaag caattgctga attggatagc 300
ctgaatgaag agtcttataa agacagcact ctgatcatgc agttacttag ggacaattca 360
ctctgtggac atcggaaaaac caggagagacg aaggagagcgc tggggaggga gagaactaat 420
gtttctcttg ctttgtgatc tgttcagtgt cactctgtac cctcaacata tatcccttgt 480
gcgataaaaa aaaaanaaaa aaaaaccntc nggggggggc ccggancccn attccccct 539

```

<210> 30

<211> 568

<212> DNA

<213> Homo sapien

<400> 30

```

attccaaacc aagtagtgtc tgtcagccct cttactctg tgcacgccct atttcagtct 60
tttacatttg ttcttctagg gaatgtatgc atctctatat atattttccc tctcaaaacc 120
agaacatcaa cagtgtctgt tctgacactt cagacatccc acgcaaagcc acattgaatt 180
tttgccaaat gaaaaacaca tccacaatca agttctaaga ggggtgtcaag tggggaatat 240
taatatgtgt tattattcaa aaatttagtt tatnaaangg aancaaaacc nttgaacctt 300
ttttccnnaa aaanaaggaa aatntnntgt ngaccaaggg ncgaacctga atccnccttg 360
aaaaattggt ntctcagaaa ggaaaagcgc cctccagttc ttttacccca agaatttana 420
aaaatttggc ccaagatttt atatgttcag ttgtttatgt ntaaaaataa ctttctggat 480
tttgtggggg aggaccggaa aaggaaggga gtttatccct atgttataca ntanaaaactt 540
cccnataaaa atgccatnga tgggttga 568

```

<210> 31

<211> 315

<212> DNA

<213> Homo sapien

<220>

<223> Human sapien

<400> 31

```

aattcggcac gagcaggag ccgctagtga aaatctggca tgaaataagg actaatggcc 60
ccaaaaaagg aggtggctct aagtaaaact gggattggac agtagtgggt catctgggtcc 120
ttgccgcctg agagccccag gagacatcgg ctagagtgc catggctatg ctcccgtctg 180
gaagatgccg gcatctggcc tcccactggt ttcagctgtg tccccagtc cgtgtctttt 240
tagaatgtga atgatgataa agttgtgaaa taaaggtttc tatctagttt gtaaaaaaaa 300
aaaaaaaaaa aaaaaa 315

```

<210> 32

<211> 458

<212> DNA

<213> Homo sapien

<400> 32

```

aattcaagga actttacatt gtaagagaaa acaaaacact gcaaaagaag tgtgccgact 60
atcaataaaa tgggtgaaatc atctgcaaat gtggccaggc ttggggaaca atgatgggtgc 120
acaaaggctt agatttgctt tgtctcaaaa taaggaattt tgtagtgggtt ttcaaaaata 180
attcaacaaa gaaacaatac aaaaagtggg tagaattacc tatcacattt cccaatcttg 240
actattcaga atgctgttta tttagtgtg aggattagca cttgattgaa gattctttta 300
aaatactatc agttaaacat ttaatatgat tatgattaat gnattcatta tgctncagac 360
tgacntanga atcantaaaa ngatngtttt actctgcaa aaaaaaaaaa aacncggggg 420
ggggcccggc cccaatttcc ccttntgggg ggggggtt 458

```

<210> 33

<211> 470

<212> DNA

<213> Homo sapien

<400> 33

```

aattcttatc ttccagaggc tacaattatt ataatggaca atacttttac ctttgtctct 60
aaagatcaga ttagttttat ttgttcactt acgtgctttg attatcccct ctgaattata 120
gaccgagctt tgttgttttag cctaagagaa gatttatgta gtaatttctt ctcagggtatg 180
gaaccacggt cataactaac atgttggcca gaatagaacc actgggttaa catattttat 240
tcaccattaa gtgatcttta tcaatattct ggattagaca acaaattacc tttctgggtg 300
tttcttgtaa actatactcc tgtttgaatg ttaaactttg ttgctaaagt ttaattttta 360
gatgtttgaa tgttcagttt atgtatttga actacaataa accaaccctt tttatatata 420
aaaaaaaaaa aacntcgagg gggggcccg cccaattnn ccctataggg 470

```

<210> 34

<211> 261

<212> DNA

<213> Homo sapien

<400> 34

```

aattcgaact gtgtgtatgt cagtggaaac aaatcaaaag ccactaacat ggctgtctgt 60
ttcactggac tgtcccattt gctgggttaa aggattgggg cccaaatcct ctggcctagc 120
atctctcagt gtttgcattt cagactgtct aaatacagca tgtgacaagc tgaagaagcc 180
aaatctagca gtcatttctg atttcattat attctcccc tcttctgtgt aaaaagacaa 240
aaaacaaaaa aaaaaaaaaa a 261

```

<210> 35

<211> 309

<212> DNA

<213> Homo sapien

<400> 35

```

aattcggcac gagctggaca ccaacagtga tggtcagcta gatttctcag aatttcttaa 60
tctgattggg ggcctagcta tggcttgcca tgactccttc ctcaaggctg tcccttccca 120
gaagcggacc tgaggacccc ttggccctgg ccttcaaacc ccccccttt cttccagcc 180
tttctgtcat catctccaca gccacccat cccctgagca cactaaccac ctcatgcagg 240
ccccacctgc caatagtaat aaagcaatgt cactttttta aaacatgaaa aaaaaaaaaa 300
aaaaaaaaa 309

```

<210> 36
 <211> 243
 <212> DNA
 <213> Homo sapien

<400> 36
 aattcggntc gagctcgaat aagtttgact tgtgttttat cttaaccacc agatcattcc 60
 ttctgtagct caggagagca cccctccacc ccatttgctc gcagtatcct agaattcttg 120
 tgctctcgct gcagttccct ttgggttcca tgttttcctt gttccctccc atgcctagct 180
 ggattgcaga gttaagttta tgattatgaa ataaaaacta aataacaaaa aaaaaaaaaa 240
 aaa 243

<210> 37
 <211> 650
 <212> DNA
 <213> Homo sapien

<400> 37
 aattcggcac gagtaccatt cagcctgaat ttgctagtgt aggctctaaa tcaagtgtag 60
 ttccgtgtga acttgccctgc agaaccaggt gtgctttgaa gcaatgttgt gggacactac 120
 cacaagcccc ttctggaaaag gatgcagaaa agaccccagc agttagcatt tcttgtttag 180
 aacttagtaa caatctagag aagaagccca ggaggactaa agctgaaaac atccctgctg 240
 ttgtgataga gattaanaac atgccaaaca aacaacctga atcatctttg tgagtcttga 300
 aaaagatgtg atatttgact ttgcttttaa actgcaagag gaaaaagact ccactgaaat 360
 tctaagtttg ccaagtagtg taattgaagt ccttgctctgg tcacacagtt taattctatt 420
 tttgtaagaa cataatggga ctgcataaca gagttctata ttacaatttt gtgattatta 480
 gtacagagta cagctatgct gtgactgttt tggaaagcca gttttaacac tatgttacat 540
 ttttgnttaa agnaagttaa accttatata acntaatgac atttgatttc tggattttcc 600
 catgataaaa aattaggggg gataaataaa aatgggttact ggaatttcaa 650

<210> 38
 <211> 687
 <212> DNA
 <213> Homo sapien

<400> 38
 gaattcggca cgagatTTTT ttatttttca ttttccctt aggcataattt agtatttttc 60
 cctcaggcag atcattctga gtgtgagagt gtgtgtgcac atgttacaaa ggcaactacc 120
 atgttaataa aatattcaat ttgaaatcct tttcgggtatt tgaattgctt ttgaataatg 180
 ttttttatct ggatgtaaca ttgttgcat agctttttta ctttccaag taattgaata 240
 cattttatta cttggacttt tataaactct ttccctaccc actataaatg agacattcac 300
 agcgttcaag tttgtattaa aggaaaggat tagtttgacc ccttcttttg atgggttaatg 360
 catacatgca gttaaattcc tttatgcaaa tgtgacactg ctttactagg tcttttagtt 420
 atttatttat tttttttttt ttgnccantt natTTTTTAN nntaattnct naaacncatt 480
 attttttttn aaaaaataaa aacacnactn tttnttttta ananttaaac cttantaaat 540
 ttttcccccn aaaaaaaatc ccntaanntt ttnaattnt tgaattnaan annaantaaa 600
 ccttttttnaa aaccnngcaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa 660

aaaaaaaaa aaaaaaaaaa aaaaaaa

687

<210> 39

<211> 2549

<212> DNA

<213> Homo sapien

<400> 39

gatggtcctt tccttctgcc acggcgggat cgggcactca cccagttgca agtgcgagca 60
ctatggagta ggcgagggtc tcgagctgtg gccgtggact taggcaacag gaaattagaa 120
atatcttctg gaaagctggc cagatttgca gatggctctg ctgtagtaca gtcagggtgac 180
actgcagtaa tggtcacagc ggtcagtaaa acaaacctt ccccttcca gtttatgcct 240
ttgggtgggtg actacagaca aaaagctgct gcagcaggtg gaattccac aaactatctg 300
agaagagagg ttggtacttc tgataaagaa attctaaca gtcgaataat agatcggtca 360
attagaccgc tctttccagc tggctacttc tatgatacac aggttctgtg taatctgtta 420
gcagtagatg gtgtaaatga gcctgatgtc ctagcaatta atggcgcttc cgtagccctc 480
tcattatcag atattccttg gaatggacct gttggggcag tacgaatagg aataattgat 540
ggagaatatg ttgttaacct aacaagaaaa gaaatgtctt ctagtacttt aaatttagtg 600
gttgctggag cacctaaaag tcagattgtc atgttggaag cctctgcaga gaacatttta 660
cagcaggact tttgccatgc tatcaaagtg ggagtgaat ataccaaca aataattcag 720
ggcattcagc agttggtaaa agaaactggg gttaccaaga ggacacctca gaagttattt 780
acctcttcgc cagagattgt gaaatatact cataaacttg ctatggagag actctatgca 840
gtttttacag attacgagca tgacaaagtt tccagagatg aagctgttaa caaaataaga 900
ttagatacgg aggaacaact aaaagaaaaa tttccagaag ccatccata tgaataata 960
gaatccttca atgttggtgc aaagggaagt tttagaagta ttgttttgaa tgaatacaaa 1020
aggtgcgatg gtcgggattt gacttcactt aggaatgtaa gttgtgaggt agatatgttt 1080
aaaacccttc atggatcagc attatttcaa agaggacaaa cacagggtgct ttgtaccgtt 1140
acatttgatt cattagaatc tggatttaag tcagatcaag ttataacagc tataaatggg 1200
ataaaagata aaaatttcat gctgcactac gagtctctc cttatgcaac taatgaaatt 1260
ggcaaagtca ctggtttaaa tagaagagaa cttgggcatg gtgctcttgc tgagaaagct 1320
ttgtatcctg ttattcccag agattttcct ttcaccataa gaggttacatc tgaagtccta 1380
gagtcaaattg ggtcatcttc tatggcatct gcatgtggcg gaagtttagc attaatggat 1440
tcaggggttc caatttcac tcgtgttgca ggcgtagcaa taggattggg caccaaaacc 1500
gatcctgaga aggtgaaat agaagattat cgtttgctga cagatatttt gggaattgaa 1560
gattacaatg gtgacatgga cttcaaaata gctggcacta ataaaggaat aactgcatta 1620
caggctgata ttaaattacc tggaatacca ataaaaattg tgatggaggc tattcaacaa 1680
gcttcagtgg caaaaaagga gatattacag atcatgaaca aaactatttc aaaacctcga 1740
gcatctagaa aagaaaatgg acctgttgta gaaactgttc aggttccatt atcaaaacga 1800
gcaaaatttg ttggacctgg tggctataac ttaaaaaaac ttcaggctga aacagggtgta 1860
actattagtc aggtggatga agaaacgttt tctgtatttg caccaacacc cagtgttatg 1920
catgaggcaa gagacttcat tactgaaatc tgcaaggatg atcaggagca gcaattagaa 1980
tttggagcag tatataccgc cacaataact gaaatcagag atactgggtg aatggtaaaa 2040
ttatatccaa atatgactgc ggtactgctt cataacacac aacttgataa cgaaagatta 2100
aacatcctac tgccctagga ttagaagttg gccaaagaaat tcaggtgaaa tactttggac 2160
gtgaccagc cgatggaaga atgaggcttt ctcgaaaagt gcttcagtcg ccagctacaa 2220
ccgtggtcag aactttgaat gacagaagta gtattgtaat gggagaacct atttcacagt 2280
catcatctaa ttctcagtga tttttttttt ttaaagagaa ttctagaatt ctattttgtc 2340
taggggtgatg tgctgtagag caacatttta gtagatcttc cattgtgtag atttctatat 2400

aatataaata cattttaatt atttgtacta aaatgctcat ttacatgtgc cattttttta 2460
 attcgagtaa cccatatttg ttttaattgta tttacattat aaatcaagaa atatttatta 2520
 ttaaaagtaa gtcatttata catcttaga 2549

<210> 40

<211> 649

<212> DNA

<213> Homo sapien

<400> 40

ttgaagatta caatgggtgac atgggacttca aaatagctgg cactaataaa ggaataactg 60
 cattacaggc tgatattaaa ttacctggaa taccaataaa aattgtgatg gaggctattc 120
 aacaagcttc agtggcaaaa aaggagatat tacagatcat gaacaaaact atttcaaaac 180
 ctcgagcatc tagaaaagaa aatggacctg ttgtagaaac tggttcagggt ccattatcaa 240
 aacgagcaaa atttgttggga cctgggtggct ataacttaaa aaaacttcag gctgaaacag 300
 gtgtaactat tagtcagggtg gatgaagaaa cgttttgtat ttgcaccaac acccagtgtt 360
 atgcatgagg caagaagact tcattactga atctgcaagg atgatcagga gcagcaatta 420
 gaatttggag cagtatatac cgccacaata actgaaatca gagatactgg tgtaatggta 480
 aaattatata caaatatgac tgcgggtactg cttcataaca cacaacttga taacgaaaga 540
 ttaaaccatcc tactgcccta ggattagaag ttggccaaga aattcagggtg aaatactttg 600
 gactgtgacc cagccgatgg aagaatgagg ctttctcgaa aagtgttc 649

<210> 41

<211> 638

<212> DNA

<213> mouse

<400> 41

aatgggtgaca tggatttcaa aatagccggt acaaataaag gaataactgc attacaggct 60
 gatattaagt tacctggagt accaattaaa attataatgg aagccatcca acaagcgtca 120
 gtggcaaaaga aggagatact gcagataatg aacaaacgat ttcaaaacct cgagcatcaa 180
 gaaaagaaaa tggaccagt ttagaaaacag taaagggtcc attatcaaaa cgagcaaaat 240
 tcgttggggc tgggtggatat cacttaaaaa aactccaggc tgagacagggt gtaacaatta 300
 gtcagggttga tgaagaaacc ttctccatat ttgcaccaac acctactgca atgcatgaag 360
 caagagattt cattacagaa atttgcagag atgatcaaga gcaacaatta gaatttggag 420
 cagtttatac cgcgacaata actgaaatca gagacactgg agtgatggta aaactgtatc 480
 caaacatgac tgcagtgtctg cttcataaatt cacaacttga ccaacgaaag attaaacatc 540
 ccactgccct aggactagag gtggccaaga aattcagggtc aaatactttg gccgtgatcc 600
 agctgatgga agaattgaggc tttctcgtaa agtacttc 638

<210> 42

<211> 705

<212> PRT

<213> Homo sapien

<400> 42

Asp Gly Pro Phe Leu Leu Pro Arg Arg Asp Arg Ala Leu Thr Gln Leu

1

5

10

15

Gln Val Arg Ala Leu Trp Ser Ser Ala Gly Ser Arg Ala Val Ala Val
 20 25 30
 Asp Leu Gly Asn Arg Lys Leu Glu Ile Ser Ser Gly Lys Leu Ala Arg
 35 40 45
 Phe Ala Asp Gly Ser Ala Val Val Gln Ser Gly Asp Thr Ala Val Met
 50 55 60
 Val Thr Ala Val Ser Lys Thr Lys Pro Ser Pro Ser Gln Phe Met Pro
 65 70 75 80
 Leu Val Val Asp Tyr Arg Gln Lys Ala Ala Ala Gly Arg Ile Pro
 85 90 95
 Thr Asn Tyr Leu Arg Arg Glu Val Gly Thr Ser Asp Lys Glu Ile Leu
 100 105 110
 Thr Ser Arg Ile Ile Asp Arg Ser Ile Arg Pro Leu Phe Pro Ala Gly
 115 120 125
 Tyr Phe Tyr Asp Thr Gln Val Leu Cys Asn Leu Leu Ala Val Asp Gly
 130 135 140
 Val Asn Glu Pro Asp Val Leu Ala Ile Asn Gly Ala Ser Val Ala Leu
 145 150 155 160
 Ser Leu Ser Asp Ile Pro Trp Asn Gly Pro Val Gly Ala Val Arg Ile
 165 170 175
 Gly Ile Ile Asp Gly Glu Tyr Val Val Asn Pro Thr Arg Lys Glu Met
 180 185 190
 Ser Ser Ser Thr Leu Asn Leu Val Val Ala Gly Ala Pro Lys Ser Gln
 195 200 205
 Ile Val Met Leu Glu Ala Ser Ala Glu Asn Ile Leu Gln Gln Asp Phe
 210 215 220
 Cys His Ala Ile Lys Val Gly Val Lys Tyr Thr Gln Gln Ile Ile Gln
 225 230 235 240
 Gly Ile Gln Gln Leu Val Lys Glu Thr Gly Val Thr Lys Arg Thr Pro
 245 250 255
 Gln Lys Leu Phe Thr Pro Ser Pro Glu Ile Val Lys Tyr Thr His Lys
 260 265 270

Leu Ala Met Glu Arg Leu Tyr Ala Val Phe Thr Asp Tyr Glu His Asp
 275 280 285

Lys Val Ser Arg Asp Glu Ala Val Asn Lys Ile Arg Leu Asp Thr Glu
 290 295 300

Glu Gln Leu Lys Glu Lys Phe Pro Glu Ala Asp Pro Tyr Glu Ile Ile
 305 310 315 320

Glu Ser Phe Asn Val Val Ala Lys Glu Val Phe Arg Ser Ile Val Leu
 325 330 335

Asn Glu Tyr Lys Arg Cys Asp Gly Arg Asp Leu Thr Ser Leu Arg Asn
 340 345 350

Val Ser Cys Glu Val Asp Met Phe Lys Thr Leu His Gly Ser Ala Leu
 355 360 365

Phe Gln Arg Gly Gln Thr Gln Val Leu Cys Thr Val Thr Phe Asp Ser
 370 375 380

Leu Glu Ser Gly Ile Lys Ser Asp Gln Val Ile Thr Ala Ile Asn Gly
 385 390 395 400

Ile Lys Asp Lys Asn Phe Met Leu His Tyr Glu Phe Pro Pro Tyr Ala
 405 410 415

Thr Asn Glu Ile Gly Lys Val Thr Gly Leu Asn Arg Arg Glu Leu Gly
 420 425 430

His Gly Ala Leu Ala Glu Lys Ala Leu Tyr Pro Val Ile Pro Arg Asp
 435 440 445

Phe Pro Phe Thr Ile Arg Val Thr Ser Glu Val Leu Glu Ser Asn Gly
 450 455 460

Ser Ser Ser Met Ala Ser Ala Cys Gly Gly Ser Leu Ala Leu Met Asp
 465 470 475 480

Ser Gly Val Pro Ile Ser Ser Ala Val Ala Gly Val Ala Ile Gly Leu
 485 490 495

Val Thr Lys Thr Asp Pro Glu Lys Gly Glu Ile Glu Asp Tyr Arg Leu
 500 505 510

Leu Thr Asp Ile Leu Gly Ile Glu Asp Tyr Asn Gly Asp Met Asp Phe
 515 520 525

Lys Ile Ala Gly Thr Asn Lys Gly Ile Thr Ala Leu Gln Ala Asp Ile
 530 535 540
 Lys Leu Pro Gly Ile Pro Ile Lys Ile Val Met Glu Ala Ile Gln Gln
 545 550 555 560
 Ala Ser Val Ala Lys Lys Glu Ile Leu Gln Ile Met Asn Lys Thr Ile
 565 570 575
 Ser Lys Pro Arg Ala Ser Arg Lys Glu Asn Gly Pro Val Val Glu Thr
 580 585 590
 Val Gln Val Pro Leu Ser Lys Arg Ala Lys Phe Val Gly Pro Gly Gly
 595 600 605
 Tyr Asn Leu Lys Lys Leu Gln Ala Glu Thr Gly Val Thr Ile Ser Gln
 610 615 620
 Val Asp Glu Glu Thr Phe Ser Val Phe Ala Pro Thr Pro Ser Val Met
 625 630 635 640
 His Glu Ala Arg Asp Phe Ile Thr Glu Ile Cys Lys Asp Asp Gln Glu
 645 650 655
 Gln Gln Leu Glu Phe Gly Ala Val Tyr Thr Ala Thr Ile Thr Glu Ile
 660 665 670
 Arg Asp Thr Gly Val Met Val Lys Leu Tyr Pro Asn Met Thr Ala Val
 675 680 685
 Leu Leu His Asn Thr Gln Leu Asp Asn Glu Arg Leu Asn Ile Leu Leu
 690 695 700
 Pro
 705

<210> 43

<211> 665

<212> PRT

<213> Homo sapien

<400> 43

Met Gly Gln Glu Lys His Val Phe Thr Ile Asp Trp Ala Gly Arg Thr
 1 5 10 15

Leu Thr Leu Thr Val Asn Tyr Glu Glu Arg Leu Tyr Ala Val Gly Lys

20	25	30
Ile Pro Gly Gly Phe Ile Lys Arg Glu Gly Arg Pro Ser Glu Lys Ala		
35	40	45
Val Leu Ala Ser Arg Leu Ile Asp Arg Pro Ile Arg Pro Leu Phe Ala		
50	55	60
Asp Gly Phe Arg Asn Glu Val Gln Val Ile Ser Ile Val Met Ser Val		
65	70	75 80
Asp Gln Asn Cys Ser Ser Glu Met Ala Ala Met Phe Gly Ser Ser Leu		
85	90	95
Ala Leu Ser Val Ser Asp Ile Pro Phe Glu Gly Pro Ile Ala Gly Val		
100	105	110
Thr Val Gly Arg Ile Asp Asp Gln Phe Ile Ile Asn Pro Thr Val Asp		
115	120	125
Gln Leu Glu Lys Ser Asp Ile Asn Leu Val Val Ala Gly Thr Lys Asp		
130	135	140
Ala Ile Asn Met Val Glu Ala Gly Ala Asp Glu Val Pro Glu Glu Ile		
145	150	155 160
Met Leu Glu Ala Ile Met Phe Gly His Glu Glu Ile Lys Arg Leu Ile		
165	170	175
Ala Phe Gln Glu Glu Ile Val Ala Ala Val Gly Lys Glu Lys Ser Glu		
180	185	190
Ile Lys Leu Phe Glu Ile Asp Glu Glu Leu Asn Glu Lys Val Lys Ala		
195	200	205
Leu Ala Glu Glu Asp Leu Leu Lys Ala Ile Gln Val His Glu Lys His		
210	215	220
Ala Arg Glu Asp Ala Ile Asn Glu Val Lys Asn Ala Val Val Ala Lys		
225	230	235 240
Phe Glu Asp Glu Glu His Asp Glu Asp Thr Ile Lys Gln Val Lys Gln		
245	250	255
Ile Leu Ser Lys Leu Val Lys Asn Glu Val Arg Arg Leu Ile Thr Glu		
260	265	270
Glu Lys Val Arg Pro Asp Gly Arg Gly Val Asp Gln Ile Arg Pro Leu		

275	280	285
Ser Ser Glu Val Gly Leu Leu Pro Arg Thr His Gly Ser Gly Leu Phe		
290	295	300
Thr Arg Gly Gln Thr Gln Ala Leu Ser Val Cys Thr Leu Gly Ala Leu		
305	310	315 320
Gly Asp Val Gln Ile Leu Asp Gly Leu Gly Val Glu Glu Ser Lys Arg		
325	330	335
Phe Met His His Tyr Asn Phe Pro Gln Phe Ser Val Gly Glu Thr Gly		
340	345	350
Pro Met Arg Gly Pro Gly Arg Arg Glu Ile Gly His Gly Ala Leu Gly		
355	360	365
Glu Arg Ala Leu Glu Pro Val Ile Pro Ser Glu Lys Asp Phe Pro Tyr		
370	375	380
Thr Val Arg Leu Val Ser Glu Val Leu Glu Ser Asn Gly Ser Thr Ser		
385	390	395 400
Gln Ala Ser Ile Cys Ala Ser Thr Leu Ala Met Met Asp Ala Gly Val		
405	410	415
Pro Ile Lys Ala Pro Val Ala Gly Ile Ala Met Gly Leu Val Lys Ser		
420	425	430
Gly Glu His Tyr Thr Val Leu Thr Asp Ile Gln Gly Met Glu Asp Ala		
435	440	445
Leu Gly Asp Met Asp Phe Lys Val Ala Gly Thr Glu Lys Gly Val Thr		
450	455	460
Ala Leu Gln Met Asp Ile Lys Ile Glu Gly Leu Ser Arg Glu Ile Leu		
465	470	475 480
Glu Glu Ala Leu Gln Gln Ala Lys Lys Gly Arg Met Glu Ile Leu Asn		
485	490	495
Ser Met Leu Ala Thr Leu Ser Glu Ser Arg Lys Glu Leu Ser Arg Tyr		
500	505	510
Ala Pro Lys Ile Leu Thr Met Thr Ile Asn Pro Asp Lys Ile Arg Asp		
515	520	525
Val Ile Gly Pro Ser Gly Lys Gln Ile Asn Lys Ile Ile Glu Glu Thr		

530 535 540
 Gly Val Lys Ile Asp Ile Glu Gln Asp Gly Thr Ile Phe Ile Ser Ser
 545 550 555 560
 Thr Asp Glu Ser Gly Asn Gln Lys Ala Lys Lys Ile Ile Glu Asp Leu
 565 570 575
 Val Arg Glu Val Glu Val Gly Gln Leu Tyr Leu Gly Lys Val Lys Arg
 580 585 590
 Ile Glu Lys Phe Gly Ala Phe Val Glu Ile Phe Ser Gly Lys Asp Gly
 595 600 605
 Leu Val His Ile Ser Glu Leu Ala Leu Glu Arg Val Gly Lys Val Glu
 610 615 620
 Asp Val Val Lys Ile Gly Asp Glu Ile Leu Val Lys Val Thr Glu Ile
 625 630 635 640
 Asp Lys Gln Gly Arg Val Asn Leu Ser Arg Lys Ala Val Leu Arg Glu
 645 650 655
 Glu Lys Glu Lys Glu Glu Gln Gln Ser
 660 665

 <210> 44
 <211> 704
 <212> PRT
 <213> Homo sapien

 <400> 44
 Asp Gly Pro Phe Leu Leu Pro Arg Arg Asp Arg Ala Leu Thr Gln Leu
 1 5 10 15
 Gln Val Arg Ala Leu Trp Ser Ser Ala Gly Ser Arg Ala Val Ala Val
 20 25 30
 Asp Leu Gly Asn Arg Lys Leu Glu Ile Ser Ser Gly Lys Leu Ala Arg
 35 40 45
 Phe Ala Asp Gly Ser Ala Val Val Gln Ser Gly Asp Thr Ala Val Met
 50 55 60
 Val Thr Ala Val Ser Lys Thr Lys Pro Ser Pro Ser Gln Phe Met Pro
 65 70 75 80

Leu Val Val Asp Tyr Arg Gln Lys Ala Ala Ala Gly Arg Ile Pro
 85 90 95

Thr Asn Tyr Leu Arg Arg Glu Val Gly Thr Ser Asp Lys Glu Ile Leu
 100 105 110

Thr Ser Arg Ile Ile Asp Arg Ser Ile Arg Pro Leu Phe Pro Ala Gly
 115 120 125

Tyr Phe Tyr Asp Thr Gln Val Leu Cys Asn Leu Leu Ala Val Asp Gly
 130 135 140

Val Asn Glu Pro Asp Val Leu Ala Ile Asn Gly Ala Ser Val Ala Leu
 145 150 155 160

Ser Leu Ser Asp Ile Pro Trp Asn Gly Pro Val Gly Val Arg Ile Gly
 165 170 175

Ile Ile Asp Gly Glu Tyr Val Val Asn Pro Thr Arg Lys Glu Met Ser
 180 185 190

Ser Ser Thr Leu Asn Leu Val Val Ala Gly Ala Pro Lys Ser Gln Ile
 195 200 205

Val Met Leu Glu Ala Ser Ala Glu Asn Ile Leu Gln Gln Asp Phe Cys
 210 215 220

His Ala Ile Lys Val Gly Val Lys Tyr Thr Gln Gln Ile Ile Gln Gly
 225 230 235 240

Ile Gln Gln Leu Val Lys Glu Thr Gly Val Thr Lys Arg Thr Pro Gln
 245 250 255

Lys Leu Phe Thr Pro Ser Pro Glu Ile Val Lys Tyr Thr His Lys Leu
 260 265 270

Ala Met Glu Arg Leu Tyr Ala Val Phe Thr Asp Tyr Glu His Asp Lys
 275 280 285

Val Ser Arg Asp Glu Ala Val Asn Lys Ile Arg Leu Asp Thr Glu Glu
 290 295 300

Gln Leu Lys Glu Lys Phe Pro Glu Ala Asp Pro Tyr Glu Ile Ile Glu
 305 310 315 320

Ser Phe Asn Val Val Ala Lys Glu Val Phe Arg Ser Ile Val Leu Asn
 325 330 335

Glu Tyr Lys Arg Cys Asp Gly Arg Asp Leu Thr Ser Leu Arg Asn Val
 340 345 350

Ser Cys Glu Val Asp Met Phe Lys Thr Leu His Gly Ser Ala Leu Phe
 355 360 365

Gln Arg Gly Gln Thr Gln Val Leu Cys Thr Val Thr Phe Asp Ser Leu
 370 375 380

Glu Ser Gly Ile Lys Ser Asp Gln Val Ile Thr Ala Ile Asn Gly Ile
 385 390 395 400

Lys Asp Lys Asn Phe Met Leu His Tyr Glu Phe Pro Pro Tyr Ala Thr
 405 410 415

Asn Glu Ile Gly Lys Val Thr Gly Leu Asn Arg Arg Glu Leu Gly His
 420 425 430

Gly Ala Leu Ala Glu Lys Ala Leu Tyr Pro Val Ile Pro Arg Asp Phe
 435 440 445

Pro Phe Thr Ile Arg Val Thr Ser Glu Val Leu Glu Ser Asn Gly Ser
 450 455 460

Ser Ser Met Ala Ser Ala Cys Gly Gly Ser Leu Ala Leu Met Asp Ser
 465 470 475 480

Gly Val Pro Ile Ser Ser Ala Val Ala Gly Val Ala Ile Gly Leu Val
 485 490 495

Thr Lys Thr Asp Pro Glu Lys Gly Glu Ile Glu Asp Tyr Arg Leu Leu
 500 505 510

Thr Asp Ile Leu Gly Ile Glu Asp Tyr Asn Gly Asp Met Asp Phe Lys
 515 520 525

Ile Ala Gly Thr Asn Lys Gly Ile Thr Ala Leu Gln Ala Asp Ile Lys
 530 535 540

Leu Pro Gly Ile Pro Ile Lys Ile Val Met Glu Ala Ile Gln Gln Ala
 545 550 555 560

Ser Val Ala Lys Lys Glu Ile Leu Gln Ile Met Asn Lys Thr Ile Ser
 565 570 575

Lys Pro Arg Ala Ser Arg Lys Glu Asn Gly Pro Val Val Glu Thr Val
 580 585 590

Gln Val Pro Leu Ser Lys Arg Ala Lys Phe Val Gly Pro Gly Gly Tyr
 595 600 605
 Asn Leu Lys Lys Leu Gln Ala Glu Thr Gly Val Thr Ile Ser Gln Val
 610 615 620
 Asp Glu Glu Thr Phe Ser Val Phe Ala Pro Thr Pro Ser Val Met His
 625 630 635 640
 Glu Ala Arg Asp Phe Ile Thr Glu Ile Cys Lys Asp Asp Gln Glu Gln
 645 650 655
 Gln Leu Glu Phe Gly Ala Val Tyr Thr Ala Thr Ile Thr Glu Ile Arg
 660 665 670
 Asp Thr Gly Val Met Val Lys Leu Tyr Pro Asn Met Thr Ala Val Leu
 675 680 685
 Leu His Asn Thr Gln Leu Asp Asn Glu Arg Leu Asn Ile Leu Leu Pro
 690 695 700

<210> 45
 <211> 245
 <212> PRT
 <213> B subtilis

<400> 45
 Asp Arg Leu Gly Leu Ala Ala Gly Gly Asp Thr Ala Val Thr Ala Pro
 1 5 10 15
 Pro Phe Pro Leu Val Tyr Ala Gly Ile Pro Arg Glu Ser Lys Leu Ser
 20 25 30
 Arg Ile Asp Arg Ile Arg Pro Leu Phe Gly Gln Val Val Asp Ala Gly
 35 40 45
 Ser Ala Leu Ser Ser Asp Ile Gly Pro Val Gly Ile Asp Asn Pro Thr
 50 55 60
 Ser Asn Leu Val Val Ala Gly Lys Ile Met Glu Ala Ala Ala Ile Gly
 65 70 75 80
 Ile Val Gly Lys Lys Leu Phe Glu Leu Ala Glu Leu Glu Lys Glu Val
 85 90 95

Glu Val Arg Ile Glu Arg Asp Gly Arg Arg Ser Glu Val His Gly Ser
 100 105 110
 Leu Phe Arg Gly Gln Thr Gln Leu Thr Leu Asp Lys Phe Met His Tyr
 115 120 125
 Phe Pro Glu Gly Gly Arg Arg Glu Gly His Gly Ala Leu Glu Ala Leu
 130 135 140
 Pro Val Ile Pro Asp Phe Pro Thr Arg Ser Glu Val Leu Glu Ser Asn
 145 150 155 160
 Gly Ser Ser Ala Ser Cys Leu Ala Met Asp Gly Val Pro Ile Val Ala
 165 170 175
 Gly Ala Gly Leu Val Glu Tyr Leu Thr Asp Ile Gly Glu Asp Gly Asp
 180 185 190
 Met Asp Phe Lys Ala Gly Thr Lys Gly Thr Ala Leu Gln Asp Ile Lys
 195 200 205
 Gly Ile Glu Ala Gln Gln Ala Glu Ile Leu Met Thr Ser Arg Pro Thr
 210 215 220
 Lys Gly Pro Gly Lys Glu Thr Gly Val Ile Thr Ser Ala Ile Gln Leu
 225 230 235 240
 Gly Val Lys Leu Glu
 245

<210> 46
 <211> 47
 <212> RNA
 <213> Homo sapien

<400> 46
 uaaauuuuau auauuuauau uuuuuaaaaua uuuauuuuauu uauuuuaa

47

<210> 47
 <211> 11
 <212> RNA
 <213> Homo sapien

<400> 47
 uauuuuauuuu a

11

<210> 48
<211> 33
<212> RNA
<213> Homo sapien

<400> 48
uauuuuauuuu aaauuuuuaa uuuuauuuuu aaau 33

<210> 49
<211> 62
<212> RNA
<213> Homo sapien

<400> 49
guuuuuuauuu uauuuuauua gauggauucu cagauuuua uauuuuuuuu uuuuuuuuuu 60
uu 62

<210> 50
<211> 111
<212> RNA
<213> Homo sapien

<400> 50
auuuuacauuu gccuuuuuuu uauuucgagu aaccuauuu uguuuuauug uuuuuacauu 60
auuuuaucaag aaauuuuuuu uuuuuuuuagu aagucuuuu uacauuuuag a 111